



00270  
PATENT & TRADEMARK OFFICE  
09/743818  
16 JAN 2001

Attorney's Docket No.: GHC11USA  
JC07 Rec'd PCT/PTO

TRANSMITTAL LETTER TO THE U.S. ELECTED OFFICE  
(EO/US) - ENTRY INTO NATIONAL STAGE UNDER 35 USC 371

PCT/AU99/00580

19 July 1999

17 July 1998

International Application No.

International Filing Date

Priority Date Claimed

PROTEASE SUSCEPTIBILITY II

Title of Invention

Anthony Steven Weiss  
235 Rainbow Street  
Randwick, New South Wales 2031  
Australia  
Citizenship: Australia  
Applicant(s) for EO/US

Box PCT  
Assistant Commissioner for Patents  
Washington, DC 20231  
Attn: EO/US

Sir:

Applicant herewith submits to the United States Elected Office  
(EO/US) the following items under 35 USC 371:

- (1) This express request to immediately begin national examination procedures (35 USC 371(f)).
- (2) A copy of the cover sheet for the published International Application along with a copy of the specification as filed: 157 pages, including 9 pages of claims, 26 sheets of formal drawings, and a copy of the 2 page International Search Report.
- (3) a copy of the 3 page Request form.
- (4) a first Preliminary Amendment for entry prior to calculation of the filing fees.

Express Mail No.

EK992703325US

JC07 Rec'd PCT/PTO 16 JAN 2001

- (5) our check in the amount of \$2,022.00, covering the basic national fee as set forth in 37 CFR 1.492(a)(3) and based on the first Preliminary Amendment.
- (6) A Second Preliminary Amendment.
- (6) A 53 page paper copy of the Sequence Listing (provided in the specification).
- (7) A 3.5" computer-readable diskette.
- (8) A 1 page Statement under 37 CFR §1.821(f).

Copies of the following miscellaneous items are also enclosed:

- (8) Copy of the 3 page Demand for International Preliminary Examination.
- (9) Copy of the 5 page Written Opinion.
- (9) Copy of the 4 page International Preliminary Examination Report.

The Combined Declaration and Power of Attorney form will be filed by the appropriate deadline under 37 CFR §1.495(c)(2) with the surcharge under 37 CFR §1.492(e).

Please charge any additional fees which may be required to effect entry into the National Phase and credit any overpayment to Deposit Account No. 08-3040.

Please direct all communications concerning this application to the undersigned.

Respectfully submitted,

HOWSON AND HOWSON  
Attorneys for the Applicants

By



Cathy A. Kodroff

Registration No. 33,980  
Spring House Corporate Center  
Box 457  
Spring House, PA 19477  
Telephone: (215) 540-9210  
Telefacsimile: (215) 540-5818

09/743818

JC07 Rec'd PTO 16 JAN 2001

GHC11USA

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of ) Group Art Unit:  
Anthony Steven Weiss )  
Appln. No. ) Examiner:  
Filed: Herewith )  
For: PROTEASE SUSCEPTIBILITY II ) January 16, 2001

Assistant Commissioner for Patents  
Washington, DC 20231

**PRELIMINARY AMENDMENT**

Sir:

Please amend the above-identified patent application as follows.

**In the Claims**

Cancel claims 23-41, 57-75, and 82-86 without prejudice.

**REMARKS**

After entry of this preliminary amendment, the pending claims are claims 1-22, 42-56, and 76-81. Claims 23-41, 57-75, and 82-86 are canceled. No new matter is introduced by this preliminary amendment.

Applicants respectfully request that this preliminary amendment be entered prior to calculating the filing fees.

Express Mail No. EK992703325US

The Director of the U. S. Patent and Trademark Office is hereby authorized to charge any deficiency in any fees due with the filing of this paper or credit any overpayment in any fees to Deposit Account No. 08-3040.

Respectfully submitted,

HOWSON AND HOWSON  
Attorneys for the Applicants

By Cathy A. Kodroff  
Cathy A. Kodroff  
Registration No. 33,980  
Spring House Corporate Center  
Box 457  
Spring House, PA 19477  
Telephone: (215) 540-9210  
Telefacsimile: (215) 540-5818

09/743818

2007 RECEIVED TPO 16 JAN 2001

GHC11USA

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of ) Group Art Unit:  
Anthony Steven Weiss ) Examiner:  
Appln. No. )  
Filed: Herewith )  
For: PROTEASE SUSCEPTIBILITY II ) January 16, 2001

Assistant Commissioner for Patents  
Washington, DC 20231

**SECOND PRELIMINARY AMENDMENT**

Sir:

Please amend the above-identified patent application as follows.

In the Specification

Please renumber original Abstract page "108" to -- 76 --.

In the Claims

Amend claims 1, 42, and 81 as follows.

1. (Amended) A method for reducing or eliminating the susceptibility of a tropoelastin to proteolysis comprising mutating a sub-sequence in the tropoelastin so that the susceptibility of the tropoelastin to proteolysis is reduced or eliminated.

42. (Amended) A method for enhancing the susceptibility of a tropoelastin to proteolysis comprising inserting a sub-sequence into the tropoelastin so that the susceptibility of the tropoelastin to proteolysis is enhanced.

Express Mail No.

EK9927033545

81. (Amended) A method for enhancing the purification of a tropoelastin comprising including a peptidomimetic molecule according to any one of claims 76 to 80 in a crude tropoelastin preparation which is being subjected to purification.

Add new claim 87 as follows.

87. A pharmaceutical composition comprising a peptidomimetic molecule according to any one of claims 76 to 80 and a pharmaceutically acceptable carrier.

#### REMARKS

Upon entry of this second preliminary amendment, claims 1-22, 42-56, 76-81, and 87 are in this application. New claim 87 is supported by original claim 82 and throughout the specification. A copy of amended claims 1, 42 and 81 showing deletions is attached as Appendix A. No new matter is added by this preliminary amendment.

Applicants respectfully request consideration of the pending claims.

The Director of the U. S. Patent and Trademark Office is hereby authorized to charge any deficiency in any fees due with the filing of this paper or credit any overpayment in any fees to our Deposit Account No. 08-3040.

Respectfully submitted,

HOWSON AND HOWSON  
Attorneys for the Applicants

By Cathy A. Kodroff  
Cathy A. Kodroff  
Registration No. 33,980  
Spring House Corporate Center  
Box 457  
Spring House, PA 19477  
Telephone: (215) 540-9210  
Telefacsimile: (215) 540-5818

## APPENDIX A - AMENDED CLAIMS

1. (Amended) A method for reducing or eliminating the susceptibility of a tropoelastin [or a tropoelastin variant] to proteolysis comprising mutating a sub-sequence in the tropoelastin [or the tropoelastin variant] so that the susceptibility of the tropoelastin [or the tropoelastin variant] to proteolysis is reduced or eliminated.

42. (Amended) A method for enhancing the susceptibility of a tropoelastin [or a tropoelastin variant] to proteolysis comprising inserting a sub-sequence into the tropoelastin [or the tropoelastin variant] so that the susceptibility of the tropoelastin [or the tropoelastin variant] to proteolysis is enhanced.

81. (Amended) A method for enhancing the purification of a tropoelastin [or a tropoelastin variant] comprising including a peptidomimetic molecule according to any one of claims 76 to 80 in a crude tropoelastin [or tropoelastin variant] preparation which is being subjected to purification.

09743818-042601

PROTEASE SUSCEPTIBILITY IITECHNICAL FIELD

The present invention relates to: manipulation of the amino acid sequence of tropoelastin, particularly human tropoelastin, to modify its protease susceptibility; to tropoelastin derivatives having modified protease susceptibility; to peptidomimetic molecules which contain amino acid sequences which correspond to or incorporate the protease susceptible sequences of tropoelastin; and to uses of the tropoelastin derivatives and peptidomimetic molecules.

The invention also relates to nucleic acid molecules and genetic constructs encoding the amino acid sequences of the derivatives and peptidomimetic molecules of the invention.

BACKGROUND ART

The insoluble cross-linked elastin molecule is highly resistant to proteolytic degradation by many proteases. However, tropoelastin, the soluble precursor of elastin, is far more vulnerable to proteolysis. Attempts at purifying tropoelastin from tissues usually result in a collection of degraded products. This degradation can be decreased by using traditional inhibitors of serine proteases (Franzblau *et al.*, 1989; Rucker, 1982, Rich and Foster, 1984; Sandberg and Wolt, 1982). Specific degradation has also been noted in cell culture of smooth muscle cells which was attributed to metalloproteinases (Hayashi *et al.*, 1995). Even highly purified tropoelastin can degrade into discrete bands on prolonged storage. This observation has led to a hypothesis that mammalian tropoelastin is occasionally co-purified with an intrinsic protease which will promote its gradual breakdown (Mecham *et al.*, 1976; Mecham *et al.*, 1977; Mecham and Foster, 1977). Experiments have shown that mammalian serum contains proteases which are capable of degrading tropoelastin (Romero *et al.*, 1986). Thus, any newly-synthesized unprotected tropoelastin exposed to

09743818-042601



blood, such as in a blood vessel wall, would be rapidly degraded. Serum has also been shown to induce elastase activity in smooth muscle cells leading to degradation of elastin (Kobayashi *et al.*, 1994). Elastin peptides are known to be chemotactic and this could be a role of tropoelastin proteolysis *in vivo* (Grosso and Scott, 1993; Bisaccia *et al.*, 1994). However, proteolysis could also result in inadequate or faulty elastin fiber repair at the site of injury. Serine protease inhibitors have been shown to reduce the degradation of tropoelastin caused by serum (Romero, *et al.*, 1986). These experiments suggested that kallikrein was a candidate serum protease. Other experiments (McGowan *et al.*, 1996) proposed that plasmin was a major protease involved. Thrombin has been used to digest heterogeneous porcine tropoelastin *in vitro* (Torres *et al.*, 1976). However, none of these studies has provided indication of where the tropoelastin molecule is cut by proteases.

#### DESCRIPTION OF THE INVENTION

In purifying a defined species of recombinant human tropoelastin (Martin *et al.*, 1995) from its fusion partner the present inventor observed limited and reproducible cleavage of the tropoelastin, by thrombin. The pattern of degradation as seen on SDS-polyacrylamide gels was similar to that seen by others during purification and storage (Mecham *et al.*, 1977). The present inventor recognised the possibility that this may be because certain portions of tropoelastin are more susceptible to protease action or are more readily available to proteases because of tropoelastin's conformation in solution. A comparison of the sizes of the protease cleavage products with the amino acid sequence of tropoelastin and the consensus cleavage sites for the proteases being examined revealed that of the many sites in the tropoelastin amino acid sequence which are homologous to consensus sequences for particular

09743818-042601

proteases, few were readily digested by proteases. By mapping the sites at which digestion was taking place, susceptible regions were identified thus providing the first precise mapping of protease cleavage sites within any tropoelastin.

From the determination of these susceptible regions, tropoelastin amino acid sequences in these susceptible regions can be modified thus providing reduced tropoelastin derivatives which have a reduced or eliminated protease susceptibility under particular conditions, as compared with the protease susceptibility of tropoelastin under the same conditions.

In the specification and claims, "reduced tropoelastin derivative" means a molecule having a modification of an amino acid sequence in a susceptible region of tropoelastin, which molecule is folded in a functional conformation. "Functional conformation" is defined below. The modification of the amino acid sequence in the susceptible region causes reduced or eliminated protease susceptibility. Reduced tropoelastin derivatives may correspond to full length tropoelastin molecules, single domains of tropoelastin which are encoded by specific exons of the tropoelastin gene or peptides which are encoded by all or part of two neighbouring exons of the tropoelastin gene.

Reduced tropoelastin derivatives may be produced by mutation events including for example, single point mutation in a nucleotide sequence which cause a residue substitution in an amino acid sequence in a susceptible region, or mutation events in a nucleotide sequence which cause an amino acid insertion or deletion in an amino acid sequence in a susceptible region. Reduced tropoelastin derivatives can also be produced by mutation of tropoelastin sequences, in regions of the tropoelastin molecule which are susceptible to protease digestion, and further mutation in other regions of tropoelastin. The

09743616-042601

further mutations may or may not alter the susceptibility of the reduced tropoelastin derivative to proteases. Reduced tropoelastin derivatives which contain these mutations may be produced synthetically.

5       Reduced tropoelastin derivatives may alternatively be produced by chemical modification of amino acid side chains in the derivative which chemically modifies a susceptible region.

10       Reduced tropoelastin derivatives may in another alternative be produced by protease digestion. Thus according to the invention, a protease digestion product of tropoelastin, which, as a result of digestion, has lost an amino acid sequence which is in a susceptible region, is a reduced tropoelastin derivative.

15       Reduced tropoelastin derivatives can also be produced by modification of tropoelastin variant amino acid sequences, in regions of the tropoelastin molecule which are susceptible to protease digestion.

20       In the specification and claims, "variants of tropoelastin" or "tropoelastin variants" means molecules which retain one or more properties of the corresponding tropoelastin molecule, for example, elastin-like properties or macro-molecular binding properties. Elastin-like properties include the phenomenon of recoil after molecular distention and the ability to undergo cross -linking and coacervation. Macro-molecular binding properties include the ability to interact with other macro-molecules, for example glycosylaminoglycans. Tropoelastin variants have an amino acid sequence which is homologous to all or part of  
30       the amino acid sequence of a tropoelastin splice form. For the purposes of this description , "homology" between the amino acid sequence of a particular variant and all or part of a tropoelastin splice form connotes a likeness short of identity, indicative of a derivation of one sequence from  
35       the other. In particular, an amino acid sequence is homologous to all or part of a tropoelastin sequence if the

09743818 042501

alignment of that amino acid sequence with the relevant tropoelastin sequence reveals an identity of about 65% over any 20 amino acid stretch or over any repetitive element of the molecules shorter than 20 amino acids in length. Such a sequence comparison can be performed via known algorithms such as that of Lipman and Pearson (1985). Tropoelastin variants may contain amino acid sequence differences as compared with tropoelastin, at a region susceptible to proteolysis, which differences do not alter the protease susceptibility of the tropoelastin variant as compared with tropoelastin. An example of such an amino acid sequence difference at a susceptible region in a tropoelastin variant may be a conservative amino acid substitution.

Thus reduced tropoelastin derivatives may be produced by mutation of a tropoelastin variant amino acid sequence, including for example, single point mutations in a nucleotide sequence which causes a residue substitution in an amino acid sequence in a susceptible region of tropoelastin. The reduced tropoelastin derivatives may also be produced by mutation of a tropoelastin variant amino acid sequence, including for example mutation events in a nucleotide sequence which cause an amino acid insertion or deletion in an amino acid sequence in a susceptible region of tropoelastin. Reduced tropoelastin derivatives can be produced by mutation of tropoelastin variant sequences, in regions of the tropoelastin molecule which are susceptible to protease digestion, and further mutation in other regions of the reduced tropoelastin variant. The further mutations may or may not alter the susceptibility of the reduced tropoelastin derivative to proteases. Reduced tropoelastin derivatives which are produced by the mutation of a tropoelastin variant may be produced synthetically.

Alternatively, reduced tropoelastin derivatives may be produced by chemical modification of amino acid side chains in the derivative which chemically modifies a susceptible

09743818-042601

region.

Alternatively, reduced tropoelastin derivatives may also be produced by protease digestion of a tropoelastin variant. Thus according to the invention, a protease  
5 digestion product of a tropoelastin variant, which, as a result of digestion, has lost an amino acid sequence in a susceptible region, is a reduced tropoelastin derivative.

It is known that tropoelastin genes in nature are expressed as multiple transcripts which are distinguished  
10 by alternative splicing of the mRNA as described in, for instance, Indik *et al* (1990); Oliver *et al* (1987); Heim *et al* (1991); Raju *et al* (1987) and Yeh *et al* (1987). The methods of the present invention can also be applied to the different splice forms of tropoelastin. The skilled  
15 addressee will readily recognise that in applying the methods of the invention to various splice forms of tropoelastin, account must be taken of the presence or absence of the identified cleavage sites in the amino acid sequence of the particular splice form in question.

Human tropoelastins are described by Indik *et al*  
20 (1990) and Tassabehji *et al* (1997). Bressan *et al* (1987) describe the amino acid sequence of chick tropoelastin, while Raju *et al* (1987) describe the amino acid sequence of bovine tropoelastin and Pierce *et al* (1992) describe the  
25 amino acid sequence of rat tropoelastin. Again taking account of variations in amino acid sequence and the existence of different splice forms, the skilled addressee will recognise that the methods of the invention can be applied to tropoelastins from other species.

In a first aspect the present invention provides a  
30 method for reducing or eliminating the susceptibility of a tropoelastin or tropoelastin variant amino acid sequence to proteolysis which method comprises mutating at least one sub-sequence in the tropoelastin or tropoelastin variant  
35 amino acid sequence, to reduce or eliminate the susceptibility of the tropoelastin or tropoelastin variant

09743818 042601

to proteolysis.

In the specification and claims, a "sub-sequence" means a sequence which is capable of being cleaved (or in other words, digested) by a protease when tropoelastin or a tropoelastin variant is folded in a functional conformation. A "functional conformation" is the conformation which imparts the elastin -like properties and macro -molecular binding properties to tropoelastin. The sub-sequences correspond to the amino acid sequences in the regions of tropoelastin which are susceptible to proteolysis.

Typically, the mutation involves altering at least one or two residues in the sub-sequence so as to reduce or eliminate susceptibility. More preferably, at least one sub-sequence is mutated. More preferably the tropoelastin is human tropoelastin.

It will be recognised that mutation to remove one or more sub-sequences which are capable of being digested by a serine protease is of particular benefit when the tropoelastin or tropoelastin variant is to be exposed to serum since the major proteolytic activity of serum for tropoelastin is serine protease activity.

In one embodiment of the first aspect of the invention, the sub-sequence is capable of being digested by a serine protease and has an amino acid sequence including the sequence RAAAG, or an amino acid sequence selected from the group of sequences shown in SEQ ID NOS: 17 to 44. When the sub-sequence is an amino acid sequence selected from the group of sequences shown in SEQ ID NOS: 17 to 44, or has an amino acid sequence including RAAAG, the sub-sequence is preferably mutated by replacing arginine in the sub-sequence with alanine. Preferably, the sub-sequence is capable of being digested by thrombin and has an amino acid sequence shown in SEQ ID NOS: 8 or 9. Preferably the sub-sequence is capable of being digested by plasmin and has an amino acid sequence shown in SEQ ID NOS: 11 or 12. More

09743616-042601

preferably, the sub-sequence is capable of being digested by kallikrein. Yet more preferably, the sub-sequence is capable of being digested by kallikrein and has an amino acid sequence shown in any one of SEQ ID NOS: 9 or 10.

5       The present inventor has noted that cleavage of SHEL and SHEL826A with metalloproteinases leads to reproducible patterns with apparently preferred cleavage sites, evidenced using methods similar to those described here. Examples of metalloproteinases include gelatinases A and B,  
10   the 72kD and 92kD proteases, and matrix metallo elastase. Significantly SDS-PAGE indicates that cleavage is, at least in some obvious instances, different to the recognition sequences seen with serine proteases as described in Table 1. Using the 92 kDa metalloproteinase, a characteristic  
15   banding pattern was obtained with clear evidence of preferred, more intense bands. For example, using methods described herein for the serine proteases, N-terminal sequencing of an approximately 10 kDa band derived from SHEL revealed the sequence: LAAAKAAYGAA. Its location  
20   in SHEL is illustrated in Figure 2. Thus a preferred recognition site resides between A and L, which is N-terminally upstream of the identified sequence of this fragment. It will be recognised that mutation to the tropoelastin or a tropoelastin variant sequence to remove  
25   one or more sub-sequences which are digested by metalloproteinases is of particular benefit when the tropoelastin or tropoelastin variant is to be exposed to, for example, wound sites, locations of tissue damage and remodelling which can expose the tropoelastin or  
30   tropoelastin variant to metalloproteinases.

In another embodiment of the first aspect of the invention, the sub-sequence is capable of being digested by a metalloproteinase and has an amino acid sequence including the sequence ALAAA, or an amino acid sequence  
35   selected from the group of sequences shown in SEQ ID NOS: 45 to 70. Preferably, the sub-sequence is capable of being

09743318.042601

digested by gelatinase A or B. Preferably the sub-sequence has the amino acid sequence shown in SEQ ID NO: 13. When the sub-sequence is an amino acid sequence selected from the group of sequences shown in SEQ ID NOS: 45 to 70, or  
5 has an amino acid sequence including ALAAA, the sub-sequence is preferably mutated by replacing alanine at any position in the sub-sequence with another amino acid residue. More preferably, the alanine N-terminal to the leucine is mutated by replacing that alanine with another  
10 amino acid residue.

In a second aspect the present invention provides a reduced tropoelastin derivative exhibiting reduced or eliminated susceptibility to proteolysis in comparison with a corresponding tropoelastin or a corresponding  
15 tropoelastin variant, the reduced tropoelastin derivative characterised in that a sub-sequence of the corresponding tropoelastin or corresponding tropoelastin variant amino acid sequence is mutated in the reduced tropoelastin derivative to eliminate or reduce the susceptibility of the  
20 reduced tropoelastin derivative to proteolysis.

Typically at least one or two residues are mutated in the sub-sequence. More preferably, at least one sub-sequence is mutated. More preferably the tropoelastin is human tropoelastin.

25 In one embodiment of the second aspect of the invention, the mutated sub-sequence has reduced or eliminated susceptibility to digestion by a serine protease. Preferably the mutated sub-sequence includes the sequence RAAAG, or is a sequence selected from the group of  
30 sequences shown in SEQ ID NOS: 17 to 44, provided that arginine in the sequence is replaced with alanine. Preferably the mutated sub-sequence has reduced or eliminated susceptibility to digestion by thrombin, and the mutated sub-sequence has the sequence shown in SEQ ID NOS:  
35 8 or 9, provided that at least one amino acid residue in the sequence is mutated. Preferably the mutated sub-

09743818-042601



sequence has reduced or eliminated susceptibility to digestion by plasmin, and the mutated sub-sequence has the sequence shown in SEQ ID NOS: 11 or 12, provided that at least one amino acid residue in the sequence is mutated.

5 More preferably, the mutated sub-sequence has reduced or eliminated susceptibility to digestion by kallikrein. Yet more preferably, the mutated sub-sequence has reduced or eliminated susceptibility to digestion by kallikrein and the mutated sub-sequence has the sequence shown in SEQ ID  
10 NOS: 9 or 10, provided that at least one amino acid residue in the sequence is mutated.

In another embodiment of the second aspect of the invention, the mutated sub-sequence has reduced or eliminated susceptibility to digestion by a  
15 metalloproteinase. Preferably the mutated sequence includes the sequence ALAAA, or is a sequence selected from the group of sequences shown in SEQ ID NOS: 45 to 70, provided that alanine at any position in the sequence is replaced with any amino acid residue except alanine. More  
20 preferably, the mutated sub-sequence has reduced or eliminated susceptibility to digestion by gelatinase A or B. More preferably, the mutated sub-sequence has reduced or eliminated susceptibility to digestion by gelatinase B and the mutated sub-sequence has the sequence shown in SEQ  
25 ID NO: 13, provided that at least one amino acid residue in the sequence is mutated. More preferably, the alanine N-terminal to the leucine is mutated by replacing that alanine with another amino acid residue.

Reduced tropoelastin derivatives of the second aspect  
30 with mutations appropriate to their use environment can beneficially be used *in vivo* at sites where there is a risk of protease attack on tropoelastin or a variant of tropoelastin, such as in the presence of serum or wound exudate. For instance, the therapeutic use of cross-linked  
35 tropoelastin or a cross-linked tropoelastin variant in blood vessel walls would benefit since serum-induced

09743818 "042601

degradation could be reduced. Further, certain modifications should reduce the need to use protease inhibitors during purification of the reduced tropoelastin derivative and result in greater amounts of full-length material if one or more susceptible regions are modified to minimise attack by endogenous host proteases.

In a third aspect the present invention provides a method of protecting a tropoelastin or a tropoelastin variant from degradation by serum or a protease selected from the group consisting of kallikrein, thrombin, trypsin and related serine proteases, including elastase, which method comprises mutating at least one sub-sequence in the tropoelastin or tropoelastin variant amino acid sequence to reduce or eliminate the susceptibility of the tropoelastin or tropoelastin variant to proteolysis. Preferably the tropoelastin is human tropoelastin. Preferably the protease is kallikrein.

In a fourth aspect the present invention provides a method of protecting a tropoelastin or a tropoelastin variant from degradation by proteolytic attack, which method comprises mutating at least one sub-sequence in the tropoelastin or tropoelastin variant amino acid sequence to reduce or eliminate the susceptibility of the tropoelastin or tropoelastin variant to proteolysis. In one embodiment the sub-sequence is digested by a metalloproteinase.

As described above, amino acid sequences of non-human tropoelastins have been determined, including the amino acid sequences of chick tropoelastin, bovine tropoelastin and rat tropoelastin (Bressan et al. 1987, Raju et al. 1987, Pierce et al. 1992). A comparison of these non-human tropoelastin amino acid sequences with tropoelastin reveals that particular regions of tropoelastin which are susceptible to proteolysis as identified in the present invention are conserved in these non-human tropoelastins. Therefore it is likely that these particular regions in the non-human tropoelastins will be susceptible to proteolysis.

09743318-042601

The analysis of the sub-sequences described in Table 1 with non human tropoelastin or elastin sequences with the 'nr' database using 'tblastn' at the NCBI Blast facility (<http://www.ncbi.nlm.nih.gov/BLAST>) shows the following:

5

(i) human tropoelastin:

554 VPTGAGVKPKAPGVGGAF 607

bovine tropoelastin, exon 14

10 373 VPTGAGVKPKAPGGGGAF 426

mouse tropoelastin mRNA complete cds

694 VPTGTGVKAKAPGGGGAF 747

15 bovine elastin a mRNA complete cds

545 VPTGAGVKPKAQVGAGAF 598

bovine elastin b mRNA complete cds

545 VPTGAGVKPKAQVGAGAF 598

20

bovine elastin c mRNA complete cds

545 VPTGAGVKPKAQVGAGAF 598

rat tropoelastin mRNA 3' end

25 646 VPTGTGVKAKVPGGGG 693

chicken tropoelastin mRNA complete cds

572 VPTGTGIKAKGPGAG 616

30 (ii) human tropoelastin:

1664 KVAAKAQLRAAAGLGAG 1714

rat tropoelastin mRNA 3' end

1837 KAAAKAQYRAAAGLGAG 1887

35

mouse tropoelastin mRNA complete cds

05743318-042601

1795 KAAAKAQYRAAAGLGAG 1845

bovine elastin a mRNA complete cds

1649 KAAAKAQFRAAAGLPAG 1699

5

bovine elastin b mRNA complete cds

1607 KAAAKAQFRAAAGLPAG 1657

bovine elastin c mRNA complete cds

10 1547 KAAAKAQFRAAAGLPAG 1597

which demonstrates that the sub-sequences identified in  
Table 1 are highly homologous with non human tropoelastin  
or elastin sequences, supporting the proposition that  
15 taking account of sequence differences the methods of the  
invention can be applied to different tropoelastin species.

This analysis also demonstrates a consensus sequence:

AKAAAKAQN<sub>0</sub>R/AAAGLN<sub>1</sub>AGN<sub>2</sub>P

wherein N<sub>0</sub> is an aromatic or hydrophobic residue;

20

N<sub>1</sub> is P or G; and

N<sub>2</sub> is a hydrophobic residue

for the site in tropoelastin which is cleaved by kallikrein  
and thrombin. An amino acid sequence which is within the  
definition of this consensus sequence may be mutated in  
25 accordance with the methods of the invention to provide the  
derivatives of the invention which have, for example,  
reduced or eliminated susceptibility to proteolysis.

In the human tropoelastin splice form described in  
more detail herein and shown in SEQ ID NO:4, the cleavage  
30 in serum occurs between residues 515 and 516; 564 and 565;  
441 and 442; 503 and 504. Thus for this splice form the  
alteration to the sequence to influence serine protease  
susceptibility preferably involves modification of at least  
one of residues 515, 516, 564, 565, 441, 442, 503, 504, 564  
35 and 565.

Alterations to reduce susceptibility to protease

092438.042601

attack can be considered to involve removal or modification of the recognition site. An example of this modification is the replacement of lysine or arginine by an amino acid residue that is not positively charged. An example of this approach is the use of leucine to replace arginine in the sequence R/AAAGLG of Table 1 using common methods of mutagenesis such as those available commercially in kit form.

Reduced tropoelastin derivatives of the invention include:

- SHELδ26a (shown in Figure 3; SEQ ID NO: 5);
- SHELδmod (shown in Figure 4; SEQ ID NO:6);
- sequences shown in SEQ ID NOS: 71 to 74.

As the inventor has determined the regions of tropoelastin which are susceptible to proteolysis, tropoelastin can be modified by inserting a sequence which corresponds to an amino acid sequence in a susceptible region of tropoelastin, into the tropoelastin amino acid sequence, thus providing an enhanced tropoelastin derivative which has enhanced protease susceptibility under particular conditions as compared with the protease susceptibility of tropoelastin under the same conditions.

Thus, in the specification and claims, "enhanced tropoelastin derivative" means a molecule produced by inserting a sequence which corresponds to an amino acid sequence in a susceptible region of tropoelastin, into the tropoelastin amino acid sequence, which molecule is folded in a functional conformation. The insertion of the amino acid sequence which corresponds to the amino acid sequence of a susceptible region causes enhanced protease susceptibility. Enhanced tropoelastin derivatives may correspond to full length tropoelastin molecules, single domains of tropoelastin which are encoded by specific exons of the tropoelastin gene or peptides which are encoded by all or part of two neighbouring exons of the tropoelastin gene.

Insertion of the amino acid sequence into tropoelastin, may occur by, for example, splicing a peptide which has an amino acid sequence which corresponds to a susceptible region in tropoelastin, into tropoelastin.

5 Thus, enhanced tropoelastin derivatives may be produced by mutation events including a mutation in a nucleotide sequence which causes an insertion of a peptide in the tropoelastin amino acid sequence wherein the inserted peptide corresponds to an amino acid sequence in a  
10 susceptible region of tropoelastin.

Alternatively, insertion of the amino acid sequence into tropoelastin may occur by modifying an amino acid sequence in a region of tropoelastin, by residue insertion, substitution or deletion, so as to generate an amino acid  
15 sequence in that region of tropoelastin which is the same as an amino acid sequence in a susceptible region of tropoelastin. Thus, enhanced tropoelastin derivatives may be produced by mutation events including a mutation in a nucleotide sequence which causes residue insertion,  
20 substitution or deletion in a region of tropoelastin, wherein the mutation events produce at the region, an amino acid sequence which corresponds to a susceptible region of tropoelastin.

Enhanced tropoelastin derivatives which have an  
25 inserted amino acid sequence in accordance with either of the above, may be mutated further by residue insertion, substitution or deletion, or further amino acid sequence insertion. The further mutations may or may not alter the susceptibility of the enhanced tropoelastin derivative to  
30 proteases. Enhanced tropoelastin derivatives which contain these mutations may be produced synthetically.

Enhanced tropoelastin derivatives can be produced by modification of tropoelastin variant amino acid sequences, in regions of tropoelastin which are susceptible to  
35 protease digestion.

Thus, enhanced tropoelastin derivatives may be

09743818.042601

produced by mutation of a tropoelastin variant amino acid sequence including a mutation in a nucleotide sequence which causes an insertion of a peptide in the tropoelastin variant amino acid sequence wherein the inserted peptide  
5 corresponds to an amino acid sequence in a susceptible region of tropoelastin.

Alternatively, enhanced tropoelastin derivatives may be produced by mutation of a tropoelastin variant amino acid sequence including a mutation in a nucleotide sequence  
10 which causes residue insertion, substitution or deletion in a region of a tropoelastin variant amino acid sequence, wherein the mutation events produce at the region, an amino acid sequence which corresponds to a susceptible region of tropoelastin.

Enhanced tropoelastin derivatives which have an inserted amino acid sequence in accordance with either of the above, may be mutated further by residue insertion, substitution or deletion, or further amino acid sequence insertion in the tropoelastin variant amino acid sequence.  
15 The further mutations may or may not alter the susceptibility of the enhanced tropoelastin derivative to proteases. Enhanced tropoelastin derivatives which contain these mutations may be produced synthetically or by recombinant methods.  
20

As described above, the tropoelastin amino acid sequence is known to be translated in various mRNA splice forms in humans and non-human animals. Further the comparison of human and non-human tropoelastin amino acid sequences reveals amino acid homology between tropoelastin  
25 amino acid sequences. Thus, these various isoforms of human and non-human tropoelastin and the mRNA splice forms encoding them can be modified to provide the enhanced tropoelastin derivatives of the invention.  
30

In a fifth aspect the invention provides a method for  
35 enhancing the susceptibility of a tropoelastin or tropoelastin variant amino acid sequence to proteolysis,

09743318 042501

which method comprises inserting a sub-sequence into a tropoelastin or tropoelastin variant amino acid sequence to enhance the susceptibility of the tropoelastin or tropoelastin variant to proteolysis. As described above, in the specification and claims, a "sub-sequence" means a sequence which is capable of being cleaved by a protease when tropoelastin or a tropoelastin variant is folded in a functional conformation. The sub-sequences correspond to the amino acid sequences in the regions of tropoelastin which are susceptible to proteolysis. Typically, at least one sub-sequence is inserted into the tropoelastin or tropoelastin variant amino acid sequence. Preferably the tropoelastin is human tropoelastin.

In one embodiment of the fifth aspect of the invention, the inserted sub-sequence is capable of being digested by a serine protease and has an amino acid sequence including the sequence RAAAG, or an amino acid sequence selected from the group of sequences shown in SEQ ID NOS: 17 to 44. Preferably, the sub-sequence is capable of being digested by thrombin and has an amino acid sequence shown in SEQ ID NOS: 8 or 9. Preferably the sub-sequence is capable of being digested by plasmin and has an amino acid sequence shown in SEQ ID NOS: 11 or 12. More preferably, the sub-sequence is capable of being digested by kallikrein. Yet more preferably, the sub-sequence is capable of being digested by kallikrein and has an amino acid sequence shown in SEQ ID NOS: 9 or 10.

In another embodiment of the fifth aspect of the invention, the sub-sequence is capable of being digested by a metalloproteinase and has an amino acid sequence including the sequence: ALAAA, or an amino acid sequence selected from the group of sequences shown in SEQ ID NOS: 45 to 70. Preferably, the sub-sequence is capable of being digested by gelatinase A or B. Preferably the sub-sequence has the amino acid sequence shown in SEQ ID NO: 13.

In a sixth aspect the invention provides an enhanced

09743313 042601



tropeoelastin derivative exhibiting enhanced susceptibility to proteolysis in comparison with a corresponding tropeoelastin or tropeoelastin variant, the enhanced tropeoelastin derivative characterised in that a sub-  
5 sequence is inserted in the amino acid sequence of the enhanced tropeoelastin derivative to enhance the susceptibility of the enhanced tropeoelastin derivative to proteolysis. Typically, at least one sub-sequence is inserted into the tropeoelastin or tropeoelastin variant  
10 amino acid sequence. Preferably the tropeoelastin is human tropeoelastin.

In one embodiment of the sixth aspect of the invention, the inserted sub-sequence is capable of being digested by a serine protease. Preferably the inserted  
15 sub-sequence includes the sequence RAAAG, or is a sequence selected from the group of sequences shown in SEQ ID NOS: 17 to 44. Preferably the inserted sub-sequence is capable of being digested by thrombin, and the inserted sub-sequence has the sequence shown in SEQ ID NOS: 8 or 9.  
20 Preferably the inserted sub-sequence is capable of being digested by plasmin, and the inserted sub-sequence has the sequence shown in SEQ ID NOS: 11 or 12. More preferably, the inserted sub-sequence is capable of being digested by kallikrein. Yet more preferably, the inserted sub-sequence  
25 is capable of being digested by kallikrein and the inserted sub-sequence has the sequence shown in SEQ ID NOS: 9 or 10.

In another embodiment of the sixth aspect of the invention, the inserted sub-sequence is capable of being digested by a metalloproteinase. Preferably the inserted  
30 sequence includes the sequence: ALAAA, or is a sequence selected from the group of sequences shown in SEQ ID NOS: 45 to 70. More preferably, the inserted sub-sequence is capable of being digested by gelatinase A or B. More preferably, the inserted sub-sequence is capable of being  
35 digested by gelatinase B and the inserted sub-sequence has the sequence shown in SEQ ID NO: 13.

09743818 042601

The enhanced tropoelastin derivative of the sixth aspect can beneficially be used *in vivo* at sites where it is desirable to augment protease attack on the derivative. Suitable molecules for manipulation include human  
5 tropoelastin molecules. In this case, the modified tropoelastin will be of use in situations in which it is desirable to have the tropoelastin or tropoelastin variant degrade rapidly. Such situations include revealing and/or release of peptides with desirable properties, to  
10 accelerate tissue repair.

As the inventor has determined the regions of tropoelastin which are susceptible to proteolysis, the susceptibility of a polypeptide to proteolysis can be modified by inserting a sequence which corresponds to an  
15 amino acid sequence in a susceptible region of tropoelastin, into the polypeptide amino acid sequence, thus providing a polypeptide derivative which has enhanced protease susceptibility under particular conditions compared with the same polypeptide which does not contain  
20 the said inserted sequence, (the corresponding polypeptide) under the same conditions.

In the specification and claims "polypeptide derivative" means a polypeptide produced by inserting a sequence which corresponds to an amino acid sequence in a  
25 susceptible region of tropoelastin, into the polypeptide sequence. The insertion of the amino acid sequence which corresponds to the amino acid sequence of a susceptible region of tropoelastin into the polypeptide sequence, causes the enhanced protease susceptibility of the  
30 polypeptide derivative.

Insertion of the amino acid sequence into the polypeptide sequence may occur by, for example, splicing a peptide which has an amino acid sequence which corresponds to a susceptible region in tropoelastin, into the  
35 polypeptide. Thus polypeptide derivatives may be produced by mutation events including a mutation in a nucleotide

09743818 042501

sequence which causes an insertion of a peptide in the polypeptide amino acid sequence wherein the inserted peptide corresponds to an amino acid sequence in a susceptible region of tropoelastin.

5       Alternatively, insertion of the amino acid sequence into the polypeptide sequence may occur by modifying an amino acid sequence in the region of the polypeptide, by residue insertion, substitution or deletion, so as to generate an amino acid sequence in that region of the  
10       polypeptide which is the same as an amino acid sequence in a susceptible region of tropoelastin. Thus, polypeptide derivatives may be produced by mutation events including a mutation in a nucleotide sequence which causes residue  
15       insertion, substitution or deletion in a region of the polypeptide, wherein the mutation events produce at the region, an amino acid sequence which corresponds to a susceptible region of tropoelastin.

      Polypeptide derivatives which contain these mutations may be produced synthetically or by recombinant DNA  
20       methods.

      Thus in a seventh aspect the invention provides a method for enhancing the susceptibility of a polypeptide amino acid sequence to proteolysis, which method comprises inserting an amino acid sequence which corresponds to an  
25       amino acid sequence in a susceptible region of tropoelastin into the polypeptide amino acid sequence to enhance the susceptibility of the polypeptide to proteolysis. Typically at least one amino acid sequence corresponding to an amino acid sequence in a susceptible region of  
30       tropoelastin is inserted into the polypeptide amino acid sequence.

      In one embodiment the inserted sequence is capable of being digested by a protease selected from the group consisting of thrombin, kallikrein, trypsin and related  
35       serine proteases including elastase. In another embodiment, the inserted sequence is digested by

09743318 042501

metalloproteinase.

In an eighth aspect, the invention provides a polypeptide derivative exhibiting enhanced susceptibility to proteolysis in comparison with a corresponding  
5 polypeptide, the polypeptide derivative characterised in that an amino acid sequence which corresponds to an amino acid sequence in a susceptible region of tropoelastin is inserted into the polypeptide amino acid sequence to enhance the susceptibility of the polypeptide to  
10 proteolysis. Typically at least one sequence corresponding to an amino acid sequence in a susceptible region of tropoelastin is inserted into the polypeptide amino acid sequence.

In one embodiment, the inserted sequence is capable of  
15 being digested by a serine protease. Preferably the serine protease is kallikrein. In another embodiment the inserted sequence may be digested by a metalloproteinase.

As the inventor has determined the regions of tropoelastin which are susceptible to proteolysis, these  
20 regions can be used to direct the specific release of peptide domains from reduced or enhanced tropoelastin derivatives of the second and sixth aspects of the invention or the specific release of peptides from the polypeptide derivatives of the eighth aspect of the  
25 invention. Typically, amino acid sequences which correspond to the susceptible regions of tropoelastin are inserted between the derivative and the peptide domain thus providing a chimeric derivative which can be digested at the susceptible region by a specific protease to release  
30 the peptide domain from the derivative.

In the specification and claims, "chimeric derivative" means a molecule produced by linking a derivative selected from the group consisting of a reduced tropoelastin derivative, enhanced tropoelastin derivative and a  
35 polypeptide derivative, with a peptide domain via an amino acid sequence which corresponds to an amino acid sequence

09743818 "042601

in a susceptible region of tropoelastin. The amino acid sequence which corresponds to the amino acid sequence of a susceptible region of tropoelastin causes the release of the peptide domain from the derivative when the chimeric derivative is digested by a specific protease.

Chimeric derivatives may be produced by recombinant DNA techniques, including for example the construction of a nucleotide sequence which encodes the derivative, the susceptible region and the peptide domain in a single open reading frame. The chimeric derivatives may alternatively be produced synthetically or by recombinant DNA methods.

Thus in a ninth aspect, the invention provides a method for producing a chimeric derivative which method comprises linking a derivative selected from the group consisting of a reduced tropoelastin derivative, enhanced tropoelastin derivative and a polypeptide derivative, with a peptide domain via an amino acid sequence which corresponds to an amino acid sequence in a susceptible region of tropoelastin.

In one embodiment, the amino acid sequence which corresponds to an amino acid sequence in a susceptible region of tropoelastin sequence may be digested by a serine protease. Preferably the serine protease is kallikrein. In another embodiment the sequence may be digested by a metalloproteinase.

In a tenth aspect, the invention provides a chimeric derivative which comprises a derivative selected from the group consisting of a reduced tropoelastin derivative, enhanced tropoelastin derivative and a polypeptide derivative, which is linked with a peptide domain via an amino acid sequence which corresponds to an amino acid sequence in a susceptible region of tropoelastin.

In one embodiment the amino acid sequence which corresponds to an amino acid sequence in a susceptible region of tropoelastin sequence may be digested by a serine protease. Preferably the serine protease is kallikrein.

In another embodiment the sequence may be digested with metalloproteinase.

The chimeric derivatives of the invention are useful where the peptide domain has a particular biological  
5 function, including for example chemotaxis, cell proliferation or cell activation. These biological functions are effected by digestion of the chimeric derivative at the sub-sequence by a particular protease so as to release the peptide domain from the derivative  
10 domain.

The mutations in accordance with this invention may be generated by conventional site-directed or random mutagenesis. Oligonucleotide-directed mutagenesis is a further option. This method comprises:

- 15 1. synthesis of an oligonucleotide with a sequence that contains the desired nucleotide substitution (mutation);
2. hybridising the oligonucleotide to a template comprising a structural sequence encoding tropoelastin; and
- 20 3. using a DNA polymerase to extend the oligonucleotide as a primer.

Another approach which is particularly suited to situations where a synthetic polynucleotide encoding the tropoelastin is prepared from oligonucleotide blocks  
25 bounded by restriction sites, is cassette mutagenesis where entire restriction fragments are replaced.

As the inventor has identified regions of tropoelastin which are susceptible to proteolysis, it is possible to use the amino acid sequences in the susceptible regions to  
30 prepare protease inhibitor molecules which are also known as peptidomimetic molecules. In the specification and claims, "peptidomimetic molecules" means molecules which imitate a region of tropoelastin which is susceptible to proteolysis, and which therefore compete with the  
35 susceptible region for the catalytic domain in a protease. Typically the peptidomimetic molecules are peptides or

09743318-042601

peptide -like.

The peptidomimetic molecules of the invention may be structurally similar to peptides. They may include an amino acid sequence of a tropoelastin or of a variant of tropoelastin which is or includes a proteolytic site. The peptidomimetic molecules of the invention may include amino acid residues which are modified at one or more chemical groups and may be linked by non-peptide bonds. These molecules can be used in situations in which it is desirable to prevent the action of the relevant proteases.

In an eleventh aspect the present invention provides a peptide or a peptidomimetic molecule including all or part of a peptide selected from the group consisting of KAPGVGGAF, RAAAGLG, RLSPELREGD, KAAQFGLVPGV, KSAAKVAAKAQLRAA, RLSPELRE and LAAAKAAKYGAA.

The peptides of this aspect of the invention may be short peptides consisting of all or part of a sequence selected from the group consisting of KAPGVGGAF, RAAAGLG, RLSPELREGD, KAAQFGLVPGV, KSAAKVAAKAQLRAA, RLSPELRE and LAAAKAAKYGAA each in combination with upstream sequence to generate a peptide typically of the order of 15 residues although it will be understood that in some cases smaller peptides could be used and frequently larger sequences could be used. The peptides can be larger molecules containing one or more of these sequences. In addition structural analogues of these peptides are included within the scope of peptidomimetic molecules of the invention, and include for instance molecules containing modified amino acid residues.

A preferred molecule is one in which the natural cleavage site would typically be located about the centre of the peptide or peptidomimetic molecule. An example peptide is H-Ala-Ala-Lys-Ala-Gln-Leu-Arg-Ala-Ala-Gly-Leu-Gly-Ala-OH which is based on the sequence RAAAGLGA, in its context within the sequence of tropoelastin(s). A peptidomimetic form of this molecule is H-Ala-Ala-Lys-Ala-

Gln-Leu-Arg-R-Ala-Ala-Ala-Gly-Leu-Gly-Ala-OH (where R = a reduced peptide bond). Also preferred are the following retro-inverso pseudo peptides: H-D-Ala-Gly-D-Leu-Gly-D-Ala-D-Ala-D-Ala-(reduced)-D-Arg-D-Leu-D-Gln-D-Ala-D-Lys-D-Ala-D-Ala-OH and H-D-Ala-Gly-D-Leu-Gly-D-Ala-D-Ala-D-Ala-D-Arg-D-Leu-D-Gln-D-Ala-D-Lys-D-Ala-D-Ala-OH. Preferably these peptides are coupled to a substrate through the N- or C-terminus.

Also preferred are the following peptides: H-Val-Pro-Gly-Ala-Leu-Ala-Ala-Ala-OH; H-Val-Pro-Gly-Ala-(reduced)-Leu-Ala-Ala-Ala-OH and the retro-inverso pseudopeptides: H-D-Ala-D-Ala-D-Ala-D-Leu-(reduced)-D-Ala-Gly-D-Pro-D-Val-OH and H-D-Ala-D-Ala-D-Ala-D-Leu-D-Ala-Gly-D-Pro-D-Val-OH. Preferably these peptides are coupled to a substrate through the N- or C- terminus.

A further category of molecules contain one or more attached reactive groups for the covalent modification of an interacting protease leading to further inhibition of activity of the protease. The invention contemplates the use of endogenous or exogenous lysyl oxidase for attaching reactive groups. It is also recognised that there is a plethora of chemically reactive groups available as biochemical reagents, which are often utilised in the construction of chemical crosslinkers. The invention contemplates the use of endogenous or exogenous lysyl oxidase for attaching reactive groups. A subset of these may be found in the Pierce Product Catalog (1997) Chapter 7 pp133 to 154. The reactive group is placed at the ends or internal to the molecule to provide a proximity to the reacting entity.

The peptides and peptidomimetic molecules of the invention are useful in a number of different environments including in the purification of tropoelastin, as a pharmaceutical agent which can be provided in an inhalant form for protecting lung tissue from damage related to elastolytic protease attack on elastin (a major cause of



lung damage in smokers) and in any other environment in which competitive inhibition of protease active sites recognising these peptides is desirable.

5 The peptides and peptidomimetic molecules of the invention are also useful in inhibiting or controlling the local growth and metastases of cancer. In particular, the inventors recognise that the peptides and peptidomimetic molecules of the invention will be useful in competing with endogenous tropoelastin for proteases which are secreted  
10 by neoplastic cells. The secretion of these proteases is typically associated with the local growth or metastases of cancer. Thus the capacity of the peptide or peptidomimetic molecules of the invention to compete with endogenous tropoelastin for the proteases may inhibit or reduce the  
15 local growth or metastasis of the cancer. In this application, the peptides or peptidomimetic molecules of the invention may be coupled to a substrate.

In a twelfth aspect the present invention provides a method for enhancing the purification of a tropoelastin or  
20 a tropoelastin variant which method comprises including at least one peptide or peptidomimetic molecule of the eleventh aspect of the invention in the crude tropoelastin or tropoelastin variant preparation which is being subjected to purification.

25 In a thirteenth aspect the present invention provides a pharmaceutical composition comprising a derivative selected from the group consisting of a reduced tropoelastin derivative, an enhanced tropoelastin derivative, a polypeptide derivative and a chimeric  
30 derivative, or a peptide or peptidomimetic molecule of the invention together with a pharmaceutically acceptable carrier or diluent. Formulations of the derivatives or peptides or peptidomimetic molecules of the present invention are prepared in accordance with standard  
35 pharmaceutical techniques. Preferred formulations in accordance with the invention include inhalant

09743818 042604

formulations, incorporation into emulsions designed for localised use, attachment to surfaces such as a stent and injectable formulations. In addition the present inventor recognises that the compositions of the invention can be adapted for use in situations in which it is desirable to limit protease activity such as that leading to clot formation.

In an fourteenth aspect the present invention provides a nucleotide sequence encoding a derivative selected from the group consisting of a reduced tropoelastin derivative, an enhanced tropoelastin derivative, a polypeptide derivative and a chimeric derivative or a peptide or peptidomimetic molecule of the invention.

The nucleotide may be provided as a recombinant DNA molecule including vector DNA. Polynucleotides can be prepared using a combination of synthetic and cDNA techniques to form hybrid modified polynucleotide molecules. These molecules also fall within the scope of this invention.

Vectors useful in this invention include plasmids, phages and phagemids. The synthetic polynucleotides of the present invention can also be used in integrative expression systems or lytic or comparable expression systems.

Suitable vectors will generally contain origins of replication and control sequences which are derived from species compatible with the intended expression host. Typically these vectors include a promoter located upstream from the polynucleotide, together with a ribosome binding site if intended for prokaryotic expression, and a phenotypic selection gene such as one conferring antibiotic resistance or supplying an auxotrophic requirement. For production vectors, vectors which provide for enhanced stability through partitioning may be chosen. Where integrative vectors are used it is not necessary for the vector to have an origin of replication. Lytic and other

09743818 042601

comparable expression systems do not need to have those functions required for maintenance of vectors in hosts.

For *E. coli* typical vectors include pBR322, pBluescript II SK<sup>+</sup>, pGEX-2T, pTrc99A, pET series vectors, particularly pET3a and pET3d, (Studier et al., 1990) and derivatives of these vectors.

In a fifteenth aspect the present invention provides a cell containing a nucleotide sequence of the fourteenth aspect of the invention.

A preferred expression system is an *E. coli* expression system. However, the invention includes within its scope the use of other hosts capable of expressing protein from the polynucleotides designed for use in *E. coli* as well as to the use of synthetic polynucleotides suitable for use in other expression systems such as other microbial expression systems. These other expression systems include yeast, and bacterial expression systems, insect cell expression systems, and expression systems involving other eukaryotic cell lines or whole organisms.

Examples of *E. coli* hosts include *E. coli* B strain derivatives (Studier et al, 1990), NM522 (Gough and Murray, 1983) and XL1-Blue (Bullock et al, 1987).

In a sixteenth aspect the present invention provides an expression product of a cell of the fifteenth aspect of the invention encoded by a nucleotide sequence of the fourteenth aspect of the invention.

The expression products of the invention may be fused expression products which include all or part of a protein encoded by the vector in peptide linkage with the expression product. They may also include, for example, an N-terminal methionine or other additional residues which do not permanently impair the elastic properties of the product.

Typically the fusion is to the N-terminus of the desired expression product. An example of a suitable protein is glutathione S-transferase (Smith and Johnson

0974381B 042501

1988). The fused protein sequence may be chosen in order to cause the expression product to be secreted or expressed as a cell surface protein to simplify purification or expressed as a cytoplasmic protein.

5 The expressed fusion products may subsequently be treated to remove the fused protein sequences to provide free modified tropoelastin. Treatment is typically through protease treatment, or in the case of secretion removal is effected by endogenous host secretion machinery. An example  
10 of this is secretion by yeasts, including but not limited to *S. cerevisiae* and *S. pombe*.

Non-fused systems include the introduction of or use of a pre-existing methionine codon. An example of this is the use of pET3a and pET3d in *E. coli*.

15 According to a seventeenth aspect of the present invention there is provided a process for the production of an expression product of the sixteenth aspect comprising:

providing a cell of the fifteenth aspect; culturing it under conditions suitable for the expression of the product  
20 of the sixteenth aspect; and collecting the expression product.

In a eighteenth aspect the present invention provides an implant formed from one or more derivatives selected from the group consisting of a reduced tropoelastin  
25 derivative, an enhanced tropoelastin derivative, a polypeptide derivative and a chimeric derivative. Where the derivative has reduced proteolytic susceptibility the implant will be intended to be maintained *in situ* over a considerable period of time whereas when the derivative has  
30 enhanced proteolytic susceptibility the implant will be intended to be maintained *in situ* over a short period of time and indeed the rapid dissolution of the implant will be desired such as where it is desired that the implant is replaced by endogenous connective tissue.

35 Tropoelastin derivatives (ie reduced tropoelastin derivatives and enhanced tropoelastin derivatives) of the

09743818, 042601

invention can be cross-linked to form elastin or elastin-like material or can be cross-linked in conjunction with other biological or synthetic molecules to form a composite material. The cross-linking of the tropoelastin derivative  
5 can be achieved by chemical oxidation of lysine side chains using processes such as ruthenium tetroxide mediated oxidation and quinone mediated oxidation, or by using bifunctional chemical cross-linking agents such as dithiobis (succinimidylpropionate), dimethyl adipimidate or  
10 dimethyl pimelimidate and those within heterologous sites such as agents that contain UV activated cross-linking domain(s). Another alternative is the cross-linking of lysine and glutamic acid side chains.

The tropoelastin derivatives (ie reduced tropoelastin  
15 derivatives and enhanced tropoelastin derivatives) may also be enzymatically cross-linked by methods including lysyl oxidase mediated oxidation or be cross-linked using gamma irradiation. The implants are formed into the required shape by cross-linking the tropoelastin derivative in a  
20 mould which conforms to the desired shape of the implant. Where the implant is required to be used in sheet form the derivative can be cross-linked on a flat surface. Relevant methodologies are described in, for example, US 4 474 851 and US 5 250 516. The elastomeric materials may be  
25 exclusively prepared from one or more derivatives or may be composites prepared from one or more derivatives together with other materials.

#### BRIEF DESCRIPTION OF THE DRAWINGS

30 Figure 1 shows a schematic diagram illustrating the relative positions of protease sites identified by N-terminal sequencing for serum, kallikrein and thrombin. Major sites are indicated with a solid bar while minor sites are indicated with a stippled bar. Since most  
35 plasmin fragments contained the same N-terminal sequence the site of cleavage could not be identified unambiguously.

The trypsin fragments identified similarly all contained the same N-terminal sequence. Therefore, the likely regions of cleavage for plasmin and trypsin are not shown.

Figure 2 shows the nucleotide sequence and amino acid sequence of SHEL. The positions of the protease recognition sites are underlined. The amino acid of SHEL is shown in SEQ ID NO:4.

Figure 3 shows the amino acid sequence of SHEL $\delta$ 26A (bottom line) compared to the amino acid sequence of SHEL. The amino acid sequence of SHEL $\delta$ 26A is shown in SEQ ID NO:5.

Figure 4 shows the nucleotide sequence and amino acid sequence of SHEL $\delta$ mod. The amino acid sequence of SHEL $\delta$ mod is shown in SEQ ID NO:6.

Figure 5 shows 10% SDS PAGE analysis of SHEL with serum after incubation for 1,2 ,3 or 18 hours (Lanes 1 to 4). Lanes 5 and 6: peptide fragments produced by serum digestion of SHEL and SHEL $\delta$ 26A respectively, purified by butanol solubilisation. Approximate sizes of fragments produced are shown in kDa. Size markers are shown in kDa.

Figure 6 shows 8% SDS-PAGE analysis of the effect of protease inhibitors on serum degradation of SHEL. Lanes 1, 3, 5, 7 and 9: SHEL incubated with serum; lane 2: SHEL incubated with serum and 0.5mM Pefabloc SC; lane 4: SHEL incubated with serum and 5mM PMSF; lane 6: SHEL incubated with serum and EDTA; lane 8: SHEL incubated with serum and 50 mM Pefabloc PK; and lane 10: SHEL incubated with serum and 1 unit Hirudin.

Figure 7 shows 8% SDS-PAGE analysis of the effect of thrombin on SHEL and SHEL $\delta$ 26A. Increasing amounts of thrombin: lane 1 (0.01 units); lane 2 (0.05 units); lane 3 (0.10 units); lane 4 (0.15 units); lane 5 (0.20 units) and lane 6 (0.25 units) were added to SHEL. Lanes 7 and 8: effect of thrombin (1U) on degradation of SHEL and SHEL $\delta$ 26A respectively. Fragment sizes are estimated in kDa. Size

markers are shown in kDa.

Figure 8 shows 8% SDS-PAGE analysis of the effect of kallikrein on SHEL and SHEL $\delta$ 26A. Increasing concentrations of kallikrein: lane 1:  $3.0 \times 10^{-4}$ ; lane 2:  $6.0 \times 10^{-4}$ ; lane 3:  $1.5 \times 10^{-3}$  and lane 4:  $3.0 \times 10^{-3}$  were added to SHEL. Lanes 5 and 6: effect of kallikrein ( $6 \times 10^{-4}$ U) on degradation of SHEL and SHEL $\delta$ 26A respectively. Fragment sizes and size markers are shown in kDa.

Figure 9 shows 10% SDS-PAGE analysis of the effect of bovine trypsin on SHEL and SHEL $\delta$ 26A. Increasing concentrations of bovine trypsin: lane 1:  $5 \times 10^{-4}$ ; lane 2:  $1 \times 10^{-3}$ ; lane 3:  $2 \times 10^{-3}$  and lane 4:  $4 \times 10^{-3}$  were added to SHEL. Lanes 5 and 6: effect of bovine trypsin ( $2 \times 10^{-3}$ U) on SHEL and SHEL $\delta$ 26A respectively. Fragment sizes and size markers are shown in kDa.

Figure 10 shows 10% SDS-PAGE analysis of the effect of plasmin on SHEL and SHEL $\delta$ 26A. Increasing concentrations of plasmin: lane 1:  $3.7 \times 10^{-7}$ ; lane 2:  $7.4 \times 10^{-7}$ ; lane 3:  $3.7 \times 10^{-6}$ ; lane 4:  $7.4 \times 10^{-6}$ ; lane 5:  $3.7 \times 10^{-5}$ ; lane 6:  $7.4 \times 10^{-5}$  were added to SHEL. Lanes 7 and 8: effect of plasmin ( $7.4 \times 10^{-5}$ U) on SHEL and SHEL $\delta$ 26A respectively. Fragment sizes and size markers are shown in kDa.

Figure 11 shows 10% SDS-PAGE analysis of the effect of human leukocyte elastase (HLE) on SHEL and SHEL $\delta$ 26A. Increasing concentrations of HLE: lane 1:  $1.6 \times 10^{-4}$ ; lane 2:  $3.2 \times 10^{-4}$ ; lane 3:  $8.0 \times 10^{-4}$ ; lane 4:  $1.6 \times 10^{-3}$ ; lane 5:  $3.2 \times 10^{-3}$  were added to SHEL. Lanes 6 and 7: effect of HLE ( $1.6 \times 10^{-3}$ U) on SHEL and SHEL $\delta$ 26A respectively. Fragment sizes and size markers are shown in kDa.

Figure 12 shows 10% SDS-PAGE analysis of the effect of S-GAL and SPS-peptide on degradation of SHEL with A: serum, 1/2 dilution 20min; B: trypsin 20min; C: plasmin  $1.5 \times 10^{-5}$ U 20min; D: kallikrein  $15 \times 10^{-4}$ U 40min; E: thrombin 0.1U 20min and F: HLE 70min. Thrombin and kallikrein were used with a 100:1 ratio. Gels were scanned by densitometry and the

relative amount of each full-length SHEL band is shown in a histogram.

Figure 13 shows SDS-PAGE analysis of the effect of coacervation on the degradation of SHEL by proteases. SHEL was incubated in the presence (+) or absence (-) of a concentration of NaCl conducive to coacervation of SHEL at 37°C with A: kallikrein; B: thrombin; C: HLE; D: trypsin; E: plasmin and F: serum; or in the presence (+) or absence (-) of a concentration of NaCl conducive to coacervation of SHEL at 16°C with G: kallikrein; H: thrombin; I: HLE; J: trypsin; K: plasmin and L: serum.

Figure 14 shows 8% SDS-PAGE gel of the effect of thrombin cleavage of soluble cell lysate containing GST-SHEL. Increasing amounts of thrombin: lane 1: 0.001 unit; lane 2: 0.005 unit; lane 3: 0.010 unit; lane 4: 0.050 unit; lane 5: 0.100 unit; lane 6: 0.500 unit and lane 7: 1.000 unit were added to soluble cell lysate.

Figure 15 shows the construction scheme for pSHELF $\delta$ 26A. pSHELF and the aberrant pSHELF $\delta$ mod were both digested with SpeI and BssHII. BssHII cuts both plasmids twice and SpeI once resulting in three fragments. The 5424 and 946bp fragments from pSHELF and the small 338bp fragment from pSHELF $\delta$ mod were purified from agarose gels. The 5424bp fragment was CIP treated to reduce recircularisation and the three fragments ligated overnight at 16°C using DNA ligase. The final product pSHELF $\delta$ 26A contained the desired deletion of exon 26A from the SHEL gene with no other mutations.

Figure 16 shows a zymogram analysis of SHEL digested with serum (Lane 1), serum with Pefabloc SC (Lane 2) or kallikrein (Lane 3).

Figure 17 shows a zymogram analysis of gelatin digested with serum in the presence of Ca<sup>2+</sup> (Lane 1), Zn<sup>2+</sup> (Lane 2), Ca<sup>2+</sup> and Zn<sup>2+</sup> (Lane 3) and Ca<sup>2+</sup>, Zn<sup>2+</sup> and EDTA (Lane 4).

Figure 18 shows a zymogram analysis of gelatin



digested with AMPA activated gelatinase A (Lane 1), unactivated gelatinase A (Lane 2) and serum (Lane 3).

Figure 19 shows protease digestion of SHEL in solution. Lane 1, standards. Lane 2, SHEL. Lane 3, SHEL plus serum. Lane 4, SHEL plus 72kDa gelatinase. Lane 5, SHEL plus 92kDa gelatinase. Lanes 6 and 7, serum plus APMA (1hr incubation), Lanes 8 and 9, serum plus APMA (overnight incubation).

Figure 20 shows human serum kallikrein digestion of SHEL in sodium phosphate buffer, pH7.8 in the presence and absence of urea. Lane 1, standards, Lane 2, SHEL (not incubated), Lane 3, SHEL incubated with buffer (no kallikrein), Lane 4, SHEL plus kallikrein, Lane 5, SHEL plus urea in buffer (no kallikrein), Lane 6, SHEL plus kallikrein in 0.3M urea, Lane 7, SHEL plus kallikrein in 1M urea.

#### BEST METHOD OF PERFORMING THE INVENTION

The recombinant and synthetic procedures used are described in standard texts such as Sambrook et al (1989).

Purification of the tropoelastin derivatives and expression products of the invention is also performed using standard techniques with the actual sequence of steps in each instance being governed by the environment from which the molecule is to be purified. By way of example, reference is made to the purification scheme disclosed in PCT/AU93/00655.

Formulations in accordance with the invention are formulated in accordance with standard techniques.

The amount of tropoelastin derivative or peptidomimetic molecule that may be combined with a carrier or diluent to produce a single dosage form will vary depending on the situation in which the formulation is to be used and the particular mode of administration.

It will be understood also that specific doses for any particular host may be influenced by factors such as the

age, sex, weight and general health of the host as well as the particular characteristics of the modified tropoelastin being used, and how it is administered.

Injectable preparations, for example, sterile  
5 injectable aqueous or oleagenous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic  
10 parenterally acceptable diluent or solvent. Among the acceptable vehicles or solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.  
15 For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid and organic solvents find use in the preparation of injectables.

Routes of administration, dosages to be administered  
20 as well as frequency of administration are all factors which can be optimised using ordinary skill in the art.

In addition, the derivatives and expression products may be prepared as topical preparations for instance as anti-wrinkle and hand lotions using standard techniques for  
25 the preparation of such formulations.

They also may be prepared in aerosol form for, for instance, administration to a patient's lungs, or in the form of surgical implants, foods or industrial products by standard techniques.

30

#### EXAMPLES

#### MATERIALS AND METHODS

35

Reagents

0943313 042601

Hirudin, PMSF, human thrombin, human plasma kallikrein, human plasmin and human leukocyte elastase (HLE) were obtained from Sigma. Bovine trypsin and Pefabloc SC were from Boehringer-Mannheim and Pefabloc PK was from Pentapharm, Switzerland. Gelatinase A (72kDa gelatinase) and gelatinase B (92kDa gelatinase) were obtained from Boehringer Mannheim Roche Diagnostics.

SHEL was obtained by the method described in WO94/14958.

10 SHEL $\delta$ 26A can be derived from SHEL by removing the synthetic coding sequence corresponding to exon 26A. A comparison of the sequence of SHEL with that of SHEL $\delta$ 26A is provided at Figure 3. Its protein product is apparently identical to a naturally made human splice form of  
15 tropoelastin.

The Transformer Mutagenesis Kit (Clontech USA) was used with pSHELF (described in WO94/14958) in accordance with the supplied protocol to remove DNA corresponding to exon 26A. The sequence of the mutagenic primer used  
20 (manufactured by Beckman Australia) was:  
5' CGG GTT TCG GTG CTG TTC CGG GCG CGC TGG 3' which flanked either side of exon 26A by 15 bp resulting in its precise deletion. A second selection primer, which mutates a unique restriction site to another restriction site is  
25 normally used in the protocol but was not in this case since deletion of exon 26A also resulted in the deletion of a unique restriction site, PmlI. This enzyme was therefore used to digest the mutation reaction to linearise any unmutated parental plasmid and consequently to enrich for  
30 mutant plasmid in accordance with the manufacturer's instructions. The reaction mixture was used to transform competent BMH17-18 *mutS* *E. coli* defective in mismatch repair, by electroporation which was performed using a Gene Pulser apparatus (BioRad USA) according to a protocol  
35 supplied by the manufacturer. Electrocompetent cells were made according to standard protocol supplied by Clontech.

Competent cells were stored in aliquots at -80°C. After electroporation cells were grown for one hour at 37°C at 280rpm in 1ml LB. The entire transformed culture was grown overnight in 5ml LB+ampicillin. Mixed plasmid DNA containing both mutated and parental plasmids was isolated from the culture using the Qiagen Spin Plasmid isolation kit and the plasmid DNA was digested with PmlI to linearise the parental plasmid. The plasmid DNA now enriched for mutated plasmid was used to transform *E. coli* HMS174 by electroporation as described above and transformants selected on LB plates containing 75µg/ml ampicillin.

Colonies were grown overnight and plasmid mini-preparations performed in which a single colony was inoculated into 3ml LB+ampicillin media in 10ml screw-topped tubes and grown overnight with shaking at 37°C. Plasmids were extracted following the alkaline lysis protocol from Sambrook et al (1989). For HMS174 two extractions with phenol/chloroform/isoamyl alcohol were performed. Constructs were screened using PmlI and those which were insensitive to digestion were further screened by KpnI/PstI double digestion. Candidate clones were sequenced (as described herein) manually using 6F (5' GGG TGT TGG CGT TGC ACC AG 3') and 7R (5' TGC ACC TAC AAC ACC GCC CG 3') primers to confirm sequence integrity either side of the deleted region.

Automated sequencing was conducted (using either Sequi-Net (Department of Biochemistry Colorado State University USA) or by SUPAMAC (Sydney University and Prince Alfred Hospital Macromolecular Analysis Centre). DNA was applied after purification by either cesium chloride gradient or Qiagen Tip 20 (Qiagen GmbH Germany) and sequenced using the same primers as for manual sequencing.) using primers 1R (5' TGC CTT TGC CGG TTT GTA CG 3') 3F (5' TCC AGG TGG CTA CGG TCT GC 3')

3R (5' GAG TAC CTA CGC CTG CGA TAC 3')

5R (5' GGA GTA CCA ACG CCG TAC TT 3')

6F (5' GGG TGT TGG CGT TGC ACC AG 3')

7R (5' TGC ACC TAC AAC ACC GCC CG 3')

5 pETforward (5' GCA CTC ACT ATA GGG AGA CC 3')

pETreverse (5' GCC AAC TCA GCT TCC TTT CG3')

was performed to verify the rest of the sequence. A number of undesired mutations were discovered necessitating further manipulation to the DNA. The mutated DNA is named

10 pSHELF $\delta$ mod.

Sequencing confirmed the region immediately surrounding the deletion was correct. PstI and BssHII restriction sites surrounding the correct region of

pSHELF $\delta$ mod was used to remove the desired segment and  
15 reinsert it into the corresponding site of pSHELF.

6.5 $\mu$ g pSHELF and 7.5 $\mu$ g pSHELF $\delta$ mod were digested with BssHII precipitated and digested with PstI. The appropriate three fragments (Figure 15) were gel purified and ligated using 1U DNA ligase (Boehringer Mannheim Germany) overnight at  
20 16°C. DNA was transformed into *E. coli* XL1-Blue and transformants selected on plates containing 75 $\mu$ g/ml ampicillin.

Plasmids were isolated by mini-preparations and screened using BglI digestion. A candidate clone was  
25 further analysed by restriction enzyme digestion and automated sequencing was then performed using primers 1R, 3F, 5R, 6F, 7R and T7 forward (5' TAA TAC GAC TCA CTA TAG GG 3') to confirm the entire sequence. The correct sequence was designated pSHELF $\delta$ 26A.

30 SHEL $\delta$ 26A displays higher protease resistance than SHEL.

#### Serum Proteolysis of SHEL

Human serum was obtained from fresh intravenous blood,  
35 centrifuged at 2000g to remove red blood cells and then

allowed to clot before serum was removed. Aliquots (20 $\mu$ l) were stored at -20°C and thawed when needed. 15 $\mu$ g tropoelastin in 50mM sodium phosphate buffer, pH 7.8 was incubated with 0.5 $\mu$ l serum in a 20 $\mu$ l reaction for between 1 and 18hr at 37°C. Similar experiments were conducted with or without the prior addition of inhibitors. Inhibitors were added at the following concentrations; 0.5 or 1Uhirudin, 0.5 or 5mM Pefabloc SC, 1 or 5mM PMSF, 25mM EDTA, 50 or 250 $\mu$ M Pefabloc PK. All inhibitors were dissolved in water except PMSF which was dissolved in isopropanol. Reactions were analysed by 8% SDS-PAGE. Serum-digested peptides to be used for sequencing were purified by the addition of 1.5 volumes n-propanol, followed by 2.5 volumes n-butanol and stirred overnight. The organic solvents were removed by rotary evaporation and peptides resuspended in 50mM sodium phosphate buffer, pH 7.8.

#### Proteolytic Assays

A range of enzyme concentrations was originally used to determine the optimal amount for subsequent experiments. Thrombin (0.01-1U), human plasma kallikrein ( $3 \times 10^{-4}$  to  $3 \times 10^{-3}$ U), human plasmin ( $7 \times 10^{-5}$  to  $4 \times 10^{-7}$ U), bovine trypsin ( $5 \times 10^{-4}$  to  $4 \times 10^{-3}$ U), and human leukocyte elastase ( $1.6 \times 10^{-4}$  to  $3.2 \times 10^{-3}$ U) were added to 10 $\mu$ g SHEL or SHEL $\delta$ 26A in 50mM sodium phosphate buffer pH 7.8 in a total volume of 20 $\mu$ l. All reactions were performed at 37°C for one hour. Gelatinase A and B were activated using 0.8mM APMA at 37°C for 30 minutes (gelatinase A) or 37°C for 45 minutes (gelatinase B). Gelatinase A ( $4 \times 10^{-3}$  -  $4 \times 10^{-2}$ ) and gelatinase B ( $2 \times 10^{-5}$  -  $1 \times 10^{-4}$ ) was added to 15mg SHEL or SHEL $\delta$ 26A in a total volume of 50mL. Gelatinase B reactions were performed in the presence of 0.75mM APMA. The degradation profile was analysed by 8, 10 or 12% SDS-PAGE.

### Zymogram analysis

8 or 10% zymogram gels were run using (1mg/ml) porcine gelatin or SHEL as the substrate. After electrophoresis, gels were washed in 2x100mL 2.5% Triton-X 100 for 20 minutes, followed by 2x100mL 50mM Tris-HCl pH7.8, 30mM NaCl for 5 minutes and incubated overnight at 37°C in 50mM Tris-HCl pH 7.8, 30mM NaCl, 5 mM CaCl<sub>2</sub>. Gels were fixed with 25% isopropanol, 10% acetic acid, washed with 3x200mL water and stained using Gelcode (Pierce).

10

### N-terminal sequencing

Gels were poured using fresh acrylamide stocks and half the usual amounts of APS and TEMED. Gels were allowed to set for 16-24hrs. For simple protein profiles, gels were pre-run at room temperature for four hours at 20mA using 150mM Tris HCl pH8.8 buffer with 10µl/L thioglycollic acid in the upper buffer chamber. Samples were loaded and run at 4°C with fresh buffer for approximately three hours. For more complex profiles gels were pre-run at room temperature in Tris-glycine buffer (25mM Tris HCl, 192mM glycine, 0.1% (w/v) SDS, pH approximately 8.3), fresh buffer added and the gel allowed to equilibrate to room temperature before samples were added and run at 20mA with 10µl/L thioglycollic acid added to the upper chamber. Pre-stained standards (Kaleidoscope; Biorad, USA) were used to monitor extent of migration.

Gels were blotted onto polyvinylidene difluoride (PVDF) membrane (ProBlott, Applied Biosystems USA) treated according to manufacturer's instructions, overnight at 70mA using 10mM CAPS pH 11.0, 10% methanol, 10 µl/L thioglycollic acid buffer at 4°C with stirring. Blotting was performed using a Hoefer Transblot apparatus and was used according to manufacturer's instructions. The membrane was stained with 0.1% Coomassie blue-R in 50% methanol and destained in 50% methanol, 10% acetic acid. The membrane was washed with water overnight before being

air-dried. Bands were excised with a clean scalpel. Samples were blotted onto PVDF as described above. Bands were excised with a clean scalpel and sequenced by Sydney University and Prince Alfred Hospital Macromolecular  
5 Analysis Centre (SUPAMAC) using Applied Biosystems hardware and protocols. Alternatively samples were sent to the Biomolecular Resource Facility Australian National University, Canberra, for sequencing.

10 Peptide Preparation and Use

S-GAL, N-VVGSPSAQDEASPLS-C, is a peptide representing the elastin binding domain of EBP (Hinek and Rabinovitch 1994). It was synthesised by Chiron Mimotopes (Australia) and purified by RP-HPLC as follows. Concentrated peptide  
15 in 50mM ammonium acetate was treated by RP-HPLC initially by perfusion chromatography (POROS, PerSeptive Biosystems USA) using an R2 reverse phase column (4.6 x 100mm) run at 9ml/min along a 0-100% acetonitrile, 0.1% trifluoroacetic acid (TFA) gradient over 7min was used. Alternatively, a  
20 Techogel10 C18 column (2.2 x 25cm) was used with a flow rate of 8ml/min. A 0-100% acetonitrile, 0.1% TFA gradient over 55 min was used after a 10min initial wash with 30% acetonitrile/0.1% TFA. The column was equilibrated for 10min between runs due to its large volume. A maximum of  
25 30-50mg peptide was loaded at any one time. For both methods sample detection was at 214 and 280nm simultaneously. Both methods were performed using Pharmacia (Sweden) pumps and detectors. The solution was removed from the collected samples by lyophilisation and  
30 purified peptide weighed to determine yield.

A large molar excess of S-GAL in Milli-Q water (10 to 200 fold) was added to 15µg SHEL in 50mM sodium phosphate pH7.8 made up to a total volume of 40µl and preincubated at  
35 (1994) before the selected protease (kallikrein,  $6-15 \times 10^{-4}$ U; thrombin 0.1-0.2U; trypsin  $2 \times 10^{-3}$ U; plasmin,  $1.5-3.7 \times 10^{-4}$ U;



<sup>5</sup>U; human leukocyte elastase,  $1.6 \times 10^{-3}$ U; serum 1 $\mu$ l) was added according to the optimal amounts determined above for 10 to 80 minutes. Various dilutions of serum from 1/2-1/50 in 50mM sodium phosphate pH7.8 were used and both SHEL and SHEL $\delta$ 26A were used for each experiment.

A peptide representing a region of SHEL cleaved by a selection of serine proteases: N-AAKAQLRAAAGLGA-C (serine protease site peptide, SPS-peptide) was synthesised by Chiron Mimotopes (Australia) to test whether its presence could protect SHEL from degradation by acting as a competitor. Experiments were conducted in parallel with S-GAL using identical procedures (see above). Both SHEL and SHEL $\delta$ 26A were used. Each reaction was analysed by 10% SDS-PAGE. Gels were scanned by densitometry and the volume of full-length SHEL calculated as follows. Scanning densitometry of stained gels was performed using the Molecular Dynamics Personal Densitometer. Images were analysed and quantitated using ImageQuant software (Version 3.2, Molecular Dynamics USA).

20

#### Proteolysis During Coacervation

10mg/ml SHEL in 50mM sodium phosphate pH7.8 and 150mM NaCl was allowed to coacervate at 37°C until cloudy before adding human plasma kallikrein ( $6 \times 10^{-4}$ U), thrombin (1U), plasmin ( $1.5 \times 10^{-5}$ U), trypsin ( $2 \times 10^{-3}$ U), HLE ( $1.6 \times 10^{-3}$ U) and serum (0.75 $\mu$ l) for one hour. Control reactions were performed at 16°C for three hours. Extent of proteolysis was monitored by SDS-PAGE.

## 30 RESULTS

### A. Degradation of SHEL by Serum

Human tropoelastin was degraded by human serum into discrete bands, resistant to further degradation. The same degradation profile was seen by SDS-PAGE with overnight

incubation as with incubations left for one hour (Figure 5). Figure 5 clearly shows the peptide fragments after purification from serum using butanol. The sizes of the major bands are approximately 50, 45, 35, 28, 27, 25, 22 and 18 kDa, visually similar to that obtained by Romero et al (1986) using porcine tropoelastin. The pattern of peptides produced was reproducible over many separate experiments. Similar results were obtained with SHELδ26A (Figure 5) but the 22 and 18kDa bands were absent and replaced by a 15kDa band.

#### B. Effect of Protease Inhibitors on Serum Degradation

Figure 6 shows the amount of full-length SHEL after incubation with serum in the presence or absence of various protease inhibitors. Wide-spectrum serine protease inhibitors were found to inhibit degradation since both Pefabloc SC and PMSF protected tropoelastin from cleavage (Figure 6). In contrast, EDTA which is an inhibitor of metalloproteinases, appeared to promote digestion. This is an unexpected result because the metalloproteinases gelatinase A and gelatinase B digest tropoelastin (Figure 19). Protease inhibitors specific for the serine proteases thrombin and kallikrein were also tested. Hirudin, a highly specific inhibitor of thrombin, did not appear to significantly inhibit degradation whereas Pefabloc PK, specific for kallikrein, inhibited proteolysis (Figure 6).

#### C. Degradation of SHEL with specific proteases

##### *Human thrombin*

Thrombin is able to cleave GST-SHEL extensively and in a reproducible manner. Cleavage of GST-SHEL bound to glutathione agarose was performed by washing and resuspending beads in 1x thrombin cleavage buffer (50mM Tris-HCl pH8.0, 150mM NaCl, 2.5mM CaCl<sub>2</sub>) and adding human thrombin (Sigma) from 0.1 to 1% (w/v) thrombin:fusion protein at 25°C for one hour (Smith and Johnston 1988).

Soluble bacterial lysates used as substrate were incubated similarly with 1x thrombin cleavage buffer, added from a 10x stock. GST (26kDa) was evident on beads by SDS-PAGE but SHEL could not be identified in the supernatant in numerous experiments. To determine whether thrombin was degrading SHEL, the entire cell lysate was subject to cleavage with increasing concentrations of thrombin. 0.01U thrombin was the lower limit for cleavage but 0.05U and greater are more effective (Figure 14). GST was clearly present. However, with 0.01U thrombin a band at approximately 64kDa could be discerned which may represent SHEL although this was not nearly as intense as the GST band. With higher thrombin concentrations this band disappeared and smaller fragments at 45, 34 and 22kDa were noted indicating that SHEL was indeed being cleaved by thrombin.

When increasing amounts of thrombin were added to pure SHEL, four major fragments were identified by SDS-PAGE estimated at 45, 34, 22 and 13 kDa (Figure 7) in addition to faint minor bands. The sizes of the major products were very similar to those seen with thrombin digests of GST-SHEL lysates. Even with an excess of thrombin added (1U/10µg SHEL) the smaller bands were resistant to further degradation whilst the 45kDa fragment disappeared. The pattern of degradation did not appear to be the same as the serum produced peptides. When the hirudin was added to reaction, degradation was inhibited (not shown) unlike the results seen with serum. The patterns of degradation seen with SHELδ26A was slightly different with the 22kDa fragment reduced in size to about 15 kDa consistent with the fragment not containing 26A (Figure 7).

#### *Human Plasma Kallikrein*

Like thrombin, increasing amounts of human plasma kallikrein added to SHEL resulted in specific and reproducible degradation. Three major fragments were identified by SDS-PAGE (Figure 8) estimated to be 45, 22

and 18kDa, in addition to faint minor bands. The major bands at 45kDa and 18kDa were resistant to further degradation whilst the 22kDa fragment eventually disappeared. Again, the pattern of degradation was not identical to that seen by serum. Pefabloc PK could inhibit degradation by plasma kallikrein (not shown). The pattern of degradation of SHELδ26A was somewhat different, with the 22 and 18kDa fragments missing and replaced by a 15kDa fragment (Figure 8), as was seen for serum.

10 *Bovine Trypsin*

Trypsin digestion of SHEL was very extensive, resulting in complete degradation with prolonged treatment. However, with dilute amounts of enzyme ( $4 \times 10^{-3}$ U) major bands could be identified at approximately 50, 45, 40, 38, 34, 31, 22 and 18kDa, giving an overall pattern similar to serum products (Figure 9). Indeed, at low enzyme concentrations the trypsin digest profile looked virtually identical to the serum digest profile. However, trypsin digestion was not easily reproducible due to the vigorous action of trypsin on SHEL. Similar results were obtained using SHELδ26A (Figure 9) except that the sizes of the smaller fragments below 34kDa were all reduced in size by approximately 4kDa and as for kallikrein and serum, the 22 and 18kDa fragments were replaced by a single fragment at 15kDa.

*Human Plasmin*

Using plasmin at low concentrations also gave a profile very similar to both serum and trypsin (Figure 10) while at high concentration extensive degradation occurred. Major bands could be isolated using low concentration plasmin at 55, 45, 40, 34, 28, 22 and 18kDa, similar but not identical to serum digested products. Similar results were obtained using SHELδ26A (Figure 10) except that smaller fragments below 34kDa were reduced by approximately 4kDa and the 22 and 18kDa fragments were replaced by 17 and 15kDa fragments.

*Human Leukocyte Elastase (HLE)*

HLE resulted in extensive degradation if left for extended period. Using  $1.6 \times 10^{-2}$ U numerous fragments were seen with two prominent fragments at 32 and 18kDa (Figure 11). Fragments were very difficult to isolate, however, and over digestion occurred easily. SHEL826A produced a similar profile but with a series of fragments appearing 4kDa smaller (Figure 11).

10 D. Zymogram analysis of serum and proteases

To confirm the identity of proteases involved in serum digestion of SHEL, a zymogram using SHEL as a substrate was used to analyse the digestion of SHEL by serum and specific proteases (Figure 16).

15 The SHEL zymogram digested with serum shows a distinct cleared zone at 64kDa and a much fainter second cleared zone (Figure 16). No cleared zones corresponding to the other serum proteases were detected in the serum. It is likely that this result was due to the abundance of these  
20 proteases in serum, and the degree of molecular unfolding of the protease in the zymogram.

The second cleared zone was not seen when the serine protease inhibitor PMSF was used in the analysis. This indicates that the second cleared zone corresponds to the  
25 digestion of SHEL by kallikrein. To further confirm kallikrein activity against SHEL, serum was electrophoresed through a zymogram gel containing SHEL, the gel strip containing serum was cut into approximately 3mm strips and each gel slice incubated with 30mg of SHEL in solution.  
30 The supernatant was then analysed by SDS-PAGE. A pattern identical to kallikrein was seen from the gel slice from the zymogram corresponding to the region for kallikrein (data not shown). This confirmed kallikrein activity in serum.

35 The 64kDa zone identified in the zymogram analysis of SHEL digested with serum did not correspond to any of the

serine proteases analysed. A 2 dimensional zymogram (first dimension isoelectric focusing gel) indicated that the isoelectric point of the enzyme which corresponds to the 64kDa zone was pI 5-5.5 (data not shown). A SwissProt database search combining pI and molecular weight indicated that the enzyme which corresponds to the 64kDa zone was likely to be either gelatinase A or B. A zymogram analysis of gelatin digested with gelatinase A or serum demonstrated a zone of digestion corresponding to 64kDa (Figure 18). This further confirms that the 64kDa zone observed in the zymogram analysis of SHEL digested with serum corresponds to gelatinase A. A cleared zone corresponding to gelatinase B is observed at a different location in this zymogram analysis. In a zymogram analysis of gelatin digested with serum, the 64kDa zone was not observed in the presence of EDTA, or in the absence of  $\text{CaCl}_2$ , or in the presence of  $\text{ZnCl}_2$  only (Figure 17). When  $\text{CaCl}_2$  or  $\text{ZnCl}_2$  was added to the digestion, the 64kDa zone was observed (Figure 17). These results further support the contention that the enzyme which corresponds with the 64kDa zone in the zymogram analysis of SHEL digested with serum is gelatinase A. Unactivated and APMA-activated gelatinase A and gelatinase B were analysed by gelatin zymography. A 64kDa zone was observed in the gelatin zymogram digested with unactivated gelatinase A (Figure 18). This indicated that the proteolytic activity observed at 64kDa in the serum digestion of the SHEL zymogram is mediated by the unactivated form of gelatinase A. A zone corresponding to approximately 60kDa was observed in the gelatin zymogram digested with APMA-activated gelatinase A (Figure 18).

#### E. Mapping of Protease-Susceptible Sites

The thrombin, kallikrein, plasmin, trypsin and serum-produced peptides indicated in Figures 5 to 11 by an arrow, were N-terminally sequenced and assigned to regions of SHEL. Peptides corresponded either to the N-terminus of

SHEL or to cleavage sites C-terminally adjacent to a Lys or Arg. Sequences of peptides are shown in Table 1 and the positions of the cleavage sites are indicated diagrammatically in Figure 1.

5       The actual sizes, in kDa, of the fragments shown in Table 1 were determined from the amino acid sequence and are shown in brackets. In some cases, this differed from the apparent size as determined by SDS-PAGE. Curiously, one site between residues 515 and 516 (Arg and Ala) was  
10       common to thrombin and kallikrein. In addition, this same site was also cleaved by human serum. This site was identified by sequencing to be located within 26A. The lack of a second kallikrein-produced fragment in SHEL  $\delta$ 26A is therefore consistent with this site being absent from  
15       this isoform. The other serum-produced bands, which were minor in comparison, were unique and appeared to consist of a mixture of peptides making the designation tentative. These peptides were the same size in both SHEL and SHEL $\delta$ 26A (Figure 7) indicating that they are predominantly N-  
20       terminal and that the other peptide fragment is present at a much lower level. Any significant proteolysis at these other sites in SHEL $\delta$ 26A should result in a 4kDa reduction in peptide size which was not evident. Due to the rampant degradation seen by both trypsin and plasmin, the smaller  
25       fragments were unable to be isolated in sufficient quantity for sequencing. However, the sizes of the fragments indicate that the 22 and 18kDa fragments of trypsin and plasmin are probably the same sequence as for kallikrein and serum. Each of the plasmin-produced bands sequenced  
30       were a mixture of the same identified sequences, not seen with any other protease or serum, and N-terminal sequence also. Since not all the plasmin and trypsin-produced peptides were able to be identified unambiguously, the likely region of cleavage for these enzymes is not shown in  
35       Figure 1.

#### F. Effect of S-Gal and SPS-peptide on Degradation

The major serine protease site (R/AAAGLG) identified in SHEL as common to thrombin, kallikrein, serum and probably trypsin and plasmin, was produced with some flanking amino acid residues as a 14 amino acid peptide (SPS-peptide). This was added to proteolytic digests of SHEL and SHEL $\delta$ 26A to assess whether this peptide could inhibit degradation by acting as an alternative site for recognition and cleavage by proteases. In addition, S-GAL, a 15 amino acid peptide corresponding to the elastin binding domain of EBP was produced to assess whether its inhibition of porcine pancreatic elastase (Hinek and Rabinovitch 1994) could be extended to other proteases with tropoelastin-degrading ability. Using a 100:1 molar excess of SPS-peptide to SHEL, more full-length SHEL was evident compared with controls using trypsin, plasmin, kallikrein and serum, judged visually by SDS-PAGE and confirmed by scanning densitometry (Figure 12). The effect was most obvious with short incubations (20 minutes) and was seen with both SHEL and SHEL $\delta$ 26A (not shown). SPS-peptide also resulted in more full-length SHEL using thrombin and HLE but to a lesser extent (Figure 12) but longer incubations with thrombin did appear to show some inhibition (Figure 12). Degradation by HLE, however, was consistently inhibited by S-GAL even with longer incubations when inhibition with SPS-peptide was no longer seen, but was not repressed altogether (Figure 12).

#### G. Effect of coacervation on degradation of SHEL

SHEL, when in the coacervated state at 37°C was significantly protected from degradation by both thrombin and kallikrein (Figure 13) but not by plasmin. There was also some inhibition of HLE, trypsin and serum (Figure 13). This inhibition of degradation was not due to the presence of high concentrations of NaCl in the reaction mixture as control reactions using both lesser concentrations of SHEL



that did not coacervate at 37°C (not shown) and reactions performed at lower temperatures not conducive to coacervation, did not show any difference in degradation in the presence or absence of NaCl (Figure 13).

5

## DISCUSSION

### Inhibition Study of Serum Degradation of SHEL.

Human serum was able to degrade tropoelastin in a specific and reproducible manner into at least five or six major peptide fragments. The SDS-PAGE banding pattern with serum is visually similar to that of Romero et al (1986).

Various inhibitor studies confirmed the protease to be a serine protease which could be inhibited by the broad spectrum serine protease inhibitors Pefabloc SC and PMSF.

The lack of inhibition of serum digestion by EDTA suggested that metalloproteinase activity was not a major contributor to SHEL digestion. Indeed, EDTA appeared to enhance degradation by serum perhaps by modulating the action of an inhibitor of serum proteases. However, it is clear that metalloproteinases digest tropoelastin because SHEL was digested with gelatinase A and gelatinase B, as demonstrated by the SDS-PAGE and zymogram analysis of SHEL digested with these enzymes.

It is expected that metalloproteinases are a major source of proteolytic activity when tropoelastin is exposed to wound exudate. Indeed, a number of studies have demonstrated the existence of metalloproteinases in wound exudate, including MMP-2 (gelatinase A) and MMP-9 (gelatinase B) (Tarlton et al. 1997). Accordingly, the invention contemplates the modification of digestion of tropoelastin by metalloproteinases in wound exudate, by use of the methods, derivatives and peptidomimetic molecules of the invention.

Thrombin did not appear to be responsible for the majority of serum cleavage because the degradation by serum

0943818 042604

was not substantially inhibited by the thrombin-specific inhibitor hirudin, yet controls using tropoelastin and thrombin were inhibited. Pefabloc PK, specific for kallikrein inhibited degradation. Romero et al (1986) found that incubation of tropoelastin with kallikrein resulted in a somewhat similar profile to its incubation with serum. The present inhibitor studies with PefablocPK are therefore consistent with kallikrein and/or proteases with similar behaviour being involved. The inhibitor Pefabloc PK is, however, not completely specific for kallikrein. According to data supplied by the manufacturer, the inhibitor constant for plasma kallikrein is  $0.7\mu\text{mol/L}$  while the next most likely enzyme to be inhibited after kallikrein is trypsin with an inhibitor constant of  $1.3\mu\text{mol/L}$  followed by plasmin at  $10\mu\text{mol/L}$ . Thus, if present in excess Pefabloc PK may be inhibiting these enzymes also. However, the lowest concentration at which complete inhibition was seen ( $50\mu\text{M}$ ) was the manufacturer's recommended amount for inhibition of kallikrein in plasma samples.

#### Identification of Serum Proteolysis

A number of enzymes have been proposed to be responsible for the serum degradation of tropoelastin. Kallikrein (Romero et al 1986) and plasmin (McGowan et al 1996) have both been put forward as potential sources of proteolysis while a trypsin-like protease was thought to be responsible for the degradation products seen when tropoelastin was isolated from tissues (Mecham and Foster 1977). A visual comparison of SHEL degradation products from serum with the individual protease digestion products revealed only a limited similarity with thrombin and kallikrein-produced peptides while trypsin and plasmin digests appeared more similar to serum-digested peptides but only when used at low concentration. Higher concentrations and/or longer incubations completely

degraded SHEL and SHEL826A in contrast to long incubations with serum which did not change the pattern greatly.

Increasing amounts of thrombin easily degraded SHEL but only three major fragments were noted, unlike serum-produced peptides where 5-6 fragments were noted. Coupled with the observation from the inhibitor studies that the thrombin-specific inhibitor hirudin did not substantially reduce serum degradation, thrombin does not appear to be the major enzyme involved in serum proteolysis of SHEL. This was corroborated by sequencing of the peptide products which showed that although one of the two sites recognised by thrombin was likewise recognised by serum, the other site was not. This may have been a consequence of low thrombin concentration but this is unlikely since both sites are recognised to a similar extent (Figure 7).

Similarly, the profile of SHEL seen after kallikrein digestion only showed limited similarity to the serum produced profile i.e. the presence of a 45kDa fragment and two fragments around 20kDa. Sequencing of the peptides showed that both the sites recognised by kallikrein were recognised by serum. The other serum-produced fragments, however, were not seen as major products of kallikrein digestion although some other fragments were present at a very low level (Figure 8). Long incubations with kallikrein (overnight) failed to increase the intensity of other fragments nor increase to resemble serum digestion products (not shown), indicating that kallikrein was unlikely to be responsible for the additional serum-produced fragments. The sequencing data, effect of a kallikrein specific protease inhibitor and visual appearance of the digestion products by SDS-PAGE are all consistent with the involvement of kallikrein in serum digestion. However the presence of other serum peptide fragments not seen as major products of kallikrein digestion indicates that kallikrein alone is not responsible for the pattern seen in serum digests.

In contrast to thrombin and kallikrein, treatment with plasmin and trypsin resulted in extensive degradation which could completely degrade SHEL if incubated for extended periods. The degradation profile seen with plasmin was quite unlike that seen by McGowan et al (1996) where only 68 and 45kDa bands were seen suggesting that the degradation had not proceeded very far in that case. Each of these digestion profiles were more similar to serum products than either thrombin or kallikrein. By visual inspection trypsin and plasmin appeared almost identical to serum digests and each other but only at a low concentration.

There was some difficulty in the sequencing of plasmin and trypsin peptides. The plasmin-produced peptides that were sequenced were found to consist of a mixture of at least two overlapping sequences at 78/79 and 81/82 (K/AAK and K/AGA) which were the same in all of the peptide fragments sequenced. In addition, sequence from the N-terminus of SHEL was also present, which made these peptides very difficult to identify unambiguously. The presence of the same peptides throughout each fragment may be an artifact resulting from this sequence co-migrating through the entire gel with other peptides and so contaminating each subsequent peptide (J. McGovern Biomolecular Resource Facility, John Curtin School of Medical Research, Australian National University, personal communication). This may have been compounded by the low levels of peptide obtained for each fragment due to the rampant degradation by plasmin.

Similarly, low levels and poor resolution made it difficult to obtain sequence for the smaller trypsin peptides. However, clear sequence data were obtained for the larger fragments which all corresponded to N-terminal sequences as was the case for the same peptides from serum. This coupled with the observation that Pefabloc PK could also inhibit trypsin in controlled reactions (not shown)

09743818 042601

and the visual similarity of peptide fragments is consistent with trypsin-like enzyme involvement with serum proteolysis but the lack of sequence data for the more informative smaller fragments means that the identification is not definitive. Similarly, the visual similarity is also consistent with plasmin involvement but this was not able to be confirmed by sequencing. Since serum proteolysis was more defined and limited than either plasmin or trypsin alone, this indicates that the presence of trypsin-like activity is probably much lower in serum and/or is more easily destroyed.

HLE digestion profile was also extensive but was different to serum, trypsin and plasmin. HLE is a serine elastase and cleaves predominantly at Val residues (Keil 1992). The difference between elastase digests of SHEL and SHEL826A was more notable as most fragments, including the largest ones, were smaller in SHEL826A, indicating that digestion was occurring preferentially from the N-terminal end which does not appear to be the case for the other enzymes or serum. HLE involvement in serum proteolysis is therefore unlikely.

Digestion with gelatinase A and gelatinase B (each previously treated with APMA) of SHEL revealed SDS-PAGE patterns of preferentially digested fragments. The banding pattern on SDS-PAGE for each of these proteases was similar, indicating that gelatinase A and gelatinase B were likely to cut at the same or identical sites. Thus the sequence specificities for these metalloproteinases were similar. These patterns differed from AMPA-treated serum, untreated serum and serine proteases. MMP-digestion revealed multiple bands. With prolonged incubation, tropoelastin displayed marked fragmentation.

In summary, by N-terminal sequencing, visual inspection of the degradation profiles by comparison with that of serum and the effect of the inhibitors the results are consistent with involvement of kallikrein and/or

0943313-042601

protease(s) capable of giving a comparable cleavage pattern, in addition to at least one other enzyme probably present at a lower level. Plasmin or another trypsin-like enzyme or combination of enzymes are the most likely to be involved in the serum digestion of SHEL. Detectable thrombin and HLE activity in serum are unlikely.

#### Mapping of Protease Sensitive Sites

The pattern of degradation of purified tropoelastin seen by others is similar to the sizes of peptides generated by our proteolysis experiments. The sizes seen by Mecham and Foster (1977) by their trypsin-like protease associated with tropoelastin, 57,45, 36, 24.5 and 13-14kDa are very similar to the number and sizes of peptides generated by serum and the individual serine proteases on both SHEL and SHEL $\delta$ 26A indicating that cleavage may be occurring in the same or similar places. A similar profile was seen with tropoelastin from human fibroblast cell culture (Davidson and Sephel 1987). Sequencing confirmed that one site between residues 515 and 516 was common to thrombin, kallikrein and serum and from the SDS-PAGE pattern, probably also plasmin and trypsin. All the peptides sequenced confirmed that cleavage occurred after a Lys or Arg as expected for many serine proteases (Keil 1992). However, tropoelastin contains a large number of Lys and Arg yet only a small number of these residues were actually recognised and cleaved. The fact that these same sites may be recognised by different serine proteases may be due to their accessibility and/or the surrounding amino acids.

Preferred recognition sites for kallikrein and thrombin are strongly influenced by the adjacent amino acid residues (Chang 1985; Keil 1992) but it would not have been possible a priori to predict where preferential cleavage occurs in human tropoelastin. For example, kallikrein cleaves preferentially at Arg residues preceded by a bulky

residue (Keil 1992). Both sites identified by N-terminal sequencing fall into this category, with Leu-Arg at 515 and Arg-Arg at 564. However, for example, another Arg preceded by a Leu at 571 does not appear to be recognised. The highly specific and limited proteolysis of SHEL and SHEL $\delta$ 26A by kallikrein has allowed kallikrein treatment to be used to produce isolated C-terminal portions of tropoelastin for further study (S. Jensen and A.S. Weiss unpublished). The thrombin sites identified, however, do not fit the preferred sites for thrombin. Thrombin recognises predominantly P2-Lys/Arg-P1' where either P2 or P1' are Gly or P4-P3-Pro-Arg/Lys-P1'-P2'', where P4 and P3 are hydrophobic and P1' and P2' are non-acidic residues (Chang 1985) with Arg greatly favoured over Lys (Keil 1992). Neither SHEL nor SHEL $\delta$ 26A contain these exact sites although the site at 152 (Lys-Pro-Lys-Ala-Pro) is similar to the latter recognition site of P3-Pro-Lys-P1'-P2'. Which sites are recognised and cleaved may therefore be under the influence of tropoelastin secondary structure. Trypsin cleaves predominantly at Arg and Lys with a preference for Arg, while plasmin preferentially cleaves at Lys (Keil 1992). Since there are more Lys than Arg in tropoelastin, it would be expected that these proteases would cleave more extensively as is shown to be the case.

25

#### Protection from Degradation

Experiments have demonstrated that EBP can protect tropoelastin from degradation by binding primarily to the VGVAPG sequence of tropoelastin (Mecham et al 1989). A peptide S-GAL which represents the elastin binding site of EBP has been used previously to model the interaction (Hinek and Rabinovitch 1994). It has been noted that S-GAL and EBP have some homology with the N-terminal sequence of proteases such as kallikrein, HLE and plasmin and are therefore proposed to bind to the same sequence in tropoelastin, thus acting as competitive inhibitors of the

proteases (Hinek and Rabinovitch 1994; Hinek et al 1993).  
Hinek and Rabinovitch (1994) showed that S-GAL could  
significantly inhibit degradation of elastin by porcine  
pancreatic elastase and inferred that HLE and other serine  
5 proteases could be similarly inhibited from degrading  
tropoelastin. In this work, the use of S-GAL did not show  
any significant or consistent inhibition of proteolysis of  
SHEL or SHEL $\delta$ 26A by serum, trypsin, plasmin or kallikrein  
although some inhibition could be seen with thrombin.  
10 However, significant and reproducible inhibition was seen  
with HLE but complete inhibition of degradation could not  
be achieved, even with the large excess of S-GAL used. The  
S-GAL used was HPLC-purified to remove any truncated  
products and it may be possible that the peptide was  
15 damaged or irreversibly denatured by this process.  
However, samples of S-GAL which were not HPLC purified gave  
similar results (not shown). The mass spectroscopy data  
supplied by the manufacturer indicated that the correct  
product was synthesised. Therefore S-GAL either did not  
20 bind to SHEL or SHEL $\delta$ 26A very effectively or was easily  
displaced by the protease. Alternatively, the proteases  
may be binding to more than one site on tropoelastin and  
are therefore not effected by S-GAL.

In summary, S-GAL showed partial inhibition of  
25 tropoelastin degradation by HLE and thrombin but  
inhibition was not as thorough as seen by Hinek and  
Rabinovitch (1994) using porcine pancreatic elastase. More  
extensive inhibition of other proteases and serum could not  
be shown consistently. N-terminal sequencing data revealed  
30 one site in SHEL which was commonly recognised by thrombin,  
kallikrein, serum and probably trypsin and plasmin. This  
site and its flanking amino acids was synthesised and this  
SPS-peptide added to the proteolytic digests of SHEL and  
SHEL $\delta$ 26A. This peptide was not expected to bind to  
35 tropoelastin but simply act as a competitor by being  
recognised by the protease thus slowing degradation of SHEL



and SHEL826A. There was reproducible evidence of protection from degradation of SHEL and SHEL826A by the presence of SPS-peptide. The amount of full-length protein was greater in the presence of SPS-peptide than in the presence of S-GAL or control digestions and was similar for both isoforms. This was most notable in the presence of low enzyme concentrations or shorter incubations and was most obvious with trypsin, plasmin, kallikrein and serum although protection from the other proteases was noted although at a reduced level. This indicates that each of the proteases and serum could recognise this peptide to some extent and therefore this is a potential inhibitor of proteolysis of tropoelastin.

There is no direct evidence that SPS-peptide is cleaved by any protease. However, the presence of a similar amount of a different peptide (S-GAL) did not exert the same effect. Thus the effect of SPS-peptide is probably not simply due to the non-specific presence of a peptide in the reaction. SPS-peptide is therefore likely to be interacting directly with the proteases (or tropoelastin) to exert its effect. SPS-peptide may allow full-length tropoelastin to persist longer in the presence of proteases, including human serum.

In summary, the inhibition of degradation of SHEL and SHEL826A by S-GAL was only noted significantly with HLE but more extensive protection could not be shown. However a reproducible inhibition was seen in the presence of SPS-peptide with each protease and serum, and was most notable with trypsin, kallikrein and serum. This peptide provides an alternative site for interaction with proteases and results in the persistence of full-length tropoelastin for longer periods.

#### Proteolysis of Coacervated Tropoelastin

Coacervation of SHEL and SHEL826A at 37°C resulted in

significant protection from proteolysis by kallikrein and thrombin and to a lesser extent by HLE, trypsin and serum. No protection was seen from attack by plasmin. The presence of 150mM NaCl did not appear to cause the inhibition since the same reactions performed under conditions not conducive to coacervation (16°C) were digested to a similar extent in the presence or absence of NaCl. Although it is possible that a simple change in conformation at 37°C could result in altered proteolytic susceptibility, this is unlikely since coacervated and non-coacervated SHEL both at 37°C were digested at different rates. The inhibition of proteolysis is therefore probably due to steric restriction in the coacervate. Of the enzymes tested, the activity of kallikrein was most significantly inhibited by coacervation. From the N-terminal sequencing results, kallikrein predominantly recognises only two sites in SHEL, both of which are in close proximity, and only one in SHELδ26A. The coacervation of tropoelastin appears to mask these sites making them less accessible to kallikrein. With thrombin, the inhibition was not as complete as with kallikrein. Thrombin recognises predominantly two sites in SHEL also but these are more distant from each other. The process of coacervation may mask these sites but if either site is slightly more accessible proteolysis would result and consequently allow easier access to the second site. Other proteases (HLE, trypsin, plasmin) and also serum recognise and cleave at many more sites within SHEL making efficient masking of all sites by coacervation unlikely and resulting in some sites remaining available for recognition and proteolysis to occur. Thus, these proteases are not as significantly inhibited by coacervation. These results indicate that in the extracellular matrix, coacervation of tropoelastin may serve an additional role to those already proposed by providing to a certain extent, protection from proteolysis including that caused by human serum. These

09743818 042601

results could be extended to the nascent elastic fibre where newly laid tropoelastin in the coacervate would be largely protected from extracellular proteases before cross-linking makes this protection essentially permanent.

5

Possible consequences of serum degradation of tropoelastin

It is clear from these results and those of others that serum contains factors capable of degrading tropoelastin. A number of serine proteases present in human blood have been shown here to be able to degrade tropoelastin specifically and reproducibly. Thus tropoelastin when secreted by cells into the extracellular matrix is vulnerable to extensive degradation prior to being insolubilised by lysyl oxidase and cross-linked. This is especially significant in blood vessels where damaged vessels may contain a number of these proteases during normal blood coagulation. Any tropoelastin secreted at this time and not protected, for example by EBP or by coacervation, would be fragmented. These results suggest that coacervation may indeed provide some protection from digestion as seen with the inhibition of degradation of coacervated SHEL (Figure 13). However, protection is by no means complete. It has previously been suggested that tropoelastin may be under negative feedback autoregulation and upon accumulation in the extracellular matrix may inhibit the production of elastin mRNA (Foster and Curtiss 1990). Elastin peptides produced by proteases such as elastase have been shown to produce negative feedback inhibition when added to undamaged fibroblast cultures while stimulating tropoelastin production in protease damaged cultures (Foster et al 1990). It has been suggested that serine protease mediated proteolysis of tropoelastin may be an important modulator of tropoelastin production and that plasmin may be involved in this process (McGowan et al 1996). Our results are consistent with this proposal although the specific enzyme(s) proposed

09743313 042601

differ slightly.

It is interesting to note that most of the cleavages identified in serum occur in the C-terminal half of the tropoelastin molecule and that most of the larger fragments were from the N-terminus (Figure 1, Table 1). Thus the action of proteases in serum on tropoelastin serves to degrade the C-terminal portion leaving a large N-terminal segment. These shortened molecules may not be incorporated into newly synthesised or growing elastic fibers due to the absence of the highly conserved C-terminus which is shown to be responsible for binding with microfibrillar proteins (Brown-Ausburger et al 1996; 1994). This is analogous to the case in supra-valvular aortic stenosis, where an elastin gene truncation results in tropoelastin missing the C-terminus with the result of severe aortic disease (Ewart et al 1994). Similarly, in fetal lamb ductus arteriosus a truncated tropoelastin missing the C terminus is not incorporated into the elastic fibre (Hinek and Rabinovitch 1993). The action of serum on human tropoelastin therefore results in tropoelastin molecules which may not be rendered insoluble and may persist in the extracellular matrix. Any fibers cross-linked may be aberrant due to improper alignment, resulting in a loss of elastic properties and strength. The persistence of soluble peptides may serve to inhibit further tropoelastin production by negative feedback inhibition (Foster and Curtiss 1990). At the same time peptides are chemotactic, as demonstrated by several studies (Bisaccia et al 1994; Grosso and Scott 1993) and may serve to recruit tissue repairing cells to the site of injury, accelerating repair of the wound. Chemotactic peptides may differ in efficacy from for example SHEL and SHELδ26A.

#### Conclusion

Human serum was shown to be capable of degrading SHEL and SHELδ26A into a number of discrete fragments. This activity was confirmed to be from a serine protease and the

regions of susceptibility to serum were precisely mapped by N-terminal sequencing. A number of other serine proteases were shown to be capable of degrading SHEL and SHEL $\delta$ 26A. From the pattern of degradation, use of selective inhibitors and N-terminal sequencing the protease responsible for serum degradation was consistent with a trypsin-like protease but kallikrein or kallikrein-like behaviour is also a likely contributor. Significant or consistent inhibition of proteolysis did not take place using S-GAL except with thrombin and HLE but reproducible inhibition was provided by SPS-peptide. However, the process of coacervation was shown to provide the most significant protection against proteolysis including by serum and was most notable for proteases which cleaved a limited number of sites.

Cleavage of SHEL and SHEL $\delta$ 26A with metalloproteinases to generate reproducible patterns with apparently preferred cleavage sites has also been demonstrated.

#### 20      INDUSTRIAL APPLICATION

The derivatives and expression products of the invention are of use in *inter alia* the medical, pharmaceutical, veterinary and cosmetic fields as tissue bulking agents, and agents for cellular chemotaxis, proliferation and growth inhibition, in particular of smooth muscle cells, epithelial cells, endothelial cells, fibroblasts, osteocytes, chondrocytes and platelets.

TABLE 1: N-terminal Sequences of Protease-Produced Tropoelastin Peptides

	Size (kDa)*	Sequence†	Position
thrombin	45	GGVPGAIPG	
	34	K/APGVGGAF	152/153
	22(19)	R/AAAGLG	515/516
kallikrein	45	GGVPGAIPG	
	22(19)	R/AAAGLG	515/516
	18(15)	R/SLSPELREGD	564/565
trypsin	55	GGVPGAIPG	
	45	GGVPGAIPG	
	40	GGVPGAIPG	
	34	GGVPGAIPG	
plasmin	55	GGVPGAIP	
	45	K/AAKAGAGL + GGVPGAIP	78/79
	40	K/AAKAGAGL +	78/79 +
	34	K/AGAGLGGV	81/82
	28	K/AAKAGAGL +	78/79 +
		K/AGAGLGGV	81/82
		K/AAAKAGAGL +	78/79 +
gelatinase B		K/AGAGLGGV	81/82
	10(12)	A/LAAKAAKYGAA	593/594
serum	50	GGVPGAIPGGVP	
	45	GGVPGAIPGG	
	34	GGVPGAIPGGVP	
	28 (25)	GGVPGAIPG +	441/442
	27	K/AAQFGLVPGV(?)‡	
	25 (20)	GGVPGAIPGGVPGGFYPG	503/504
	22 (19)	GGVPGAIPG + K/SAAKVAAKAQ(?)	515/516
	18 (15)	R/AAAGLG	564/565
	13	R/SLSPELRE	
		GGVPGAIP	
<p>* Size of fragments are calculated from SDS-PAGE and are approximate. Sizes in brackets are the sizes determined from the position of the cleavage determined by N-terminal sequencing.</p> <p>† A slash (/) indicates an internal cleavage site adjacent to an R or K residue (bold). N-terminal sequence of residues to the right of these sites was obtained allowing the precise location of the cleavage site to be allocated and the exact size of the fragment to be calculated.</p> <p>‡ A question mark (?) indicates that this designation is tentative. The peptide is likely to be present at a very low level and as a mixture with other peptides.</p>			

REFERENCES

1. Indik Z, Yeh H, Ornstein-Goldstein N, Sheppard P,  
Anderson N, Rosenbloom JC, Peltonen L and Rosenbloom J  
5 (1987) *PNAS* (USA) 84 5680-5684
2. Indik Z, Abrams W.R., Kucich U, Gibson C.W., Mecham  
R.P. and Rosenbloom J (1990) *Arch. Biochem Biophys* 280 80-  
86
3. Oliver L, Luvalle PA, Davidson J.M., Rosenbloom J,  
10 Mathew C.G., Betser A.J. and Boyd C.D. (1987) *Collagen Rel*  
*Res* 7 77-89
4. Sambrook J., Fritsch E.F., and Maniatis T. (1989)  
*Molecular Cloning: a laboratory Manual, second edition Cold*  
*Spring Harbor Laboratory Press, Cold Spring Harbor , New*  
15 *York*
5. Bressan G.M., Argos P. and Stanley K.K. (1987)  
*Biochemistry* 26 1497-11503
6. Raju K. and Anwar R.A. (1987) *J. Biol Chem* 262 5755-  
5762
- 20 7. Pierce R.A., Alatawi A, Deak S.B. & Boyd C.D. (1992)  
*Genomics* 12 651-658
8. Lipman and Pearson (1985) *Science* 227,1435.
9. Bedell-Hogan, D., Trackman, P., Abrams, W.,  
Rosenbloom, J. and Kagan H. (1993) *J. Biol. Chem.* 268,  
25 10345-10350
10. Studier, F. W., Rosenberg, A. H., Dunn, J. J. and  
Dubendorff, J. W. (1990) *Methods Enzymol.* 185, 60-89
11. Gough, J., and Murray, N. (1983) *J. Mol. Biol.* 166,  
1-19
- 30 12. Bullock, W. O., Fernandez, J. M. and Short, J. M.  
(1987) *BioTechniques* 5, 376-379
13. Bisaccia F, Catiglione Morelli M A, De Biasi M,  
Traniello S, and Tarnburro A M. (1994)  
*Int. J. Peptide Protein Res*, 44: 332-341
- 35 14. Chang J Y (1985) *Eur. J. Biochem.* 151: 217-224

15. Grosso L E and Scott M (1993)  
Arch. Biochem. Biophys. 305: 401-404
16. Franzblau C, Pratt C A, Faris B, Colannino N M, Offner  
G D, Mogayzel P J and Troxler R F (1989)  
5 J. Biol. Chem. 264: 15115-15119
17. Hayashi A, Wachi M and Tajima S (1995)  
Biochim. Biophys. Acta 1244: 325-330
18. Keil B (1992) *Specificity of Proteolysis*
19. Kobayashi J, Wigle D, Childs T, Zhu, L, Keeley F W and  
10 Rabinovitch M (1994) J. Cell. Physiol. 160: 121-131
20. Martin S L, Vrhovski B and Weiss A S (1995)  
Gene 154, 159-166
21. Mecham R P, Foster J A and Franzblau C (1976)  
Biochim. Biophys. Acta. 446: 245-2,54
- 15 22. Mecham R P and Foster J A (1977)  
Biochem. 16: 3825-3831.
23. Mecham R P, Foster J A and Franzblau C (1977)  
Adv. Expt. Med. Biol. 79: 209-216
24. Rich C B and Foster J A (1984)  
20 Biochem. J. 217: 581-584
25. Romero N, Tinker D, Hyde D and Rucker R B (1986)  
Arch. Biochem. Biophys. 244: 161-168
26. Rucker R B (1982) Meth. Enzym. 82: 650-657
27. Sandberg L B and Wolt T B (1982) Meth. Enzym. 82: 657-  
25 665
28. Torres A R, Alvarez V L, Sandberg L B and Gray W R  
(1977) Adv. Expt. Med. Biol. 79: 267-276
29. McGowan S E, Lui R, Harvey C S and Jaeckel E C (1996)  
Am. J. Physiol. 270: L376-l385.
- 30 30. Heim R A, Pierce R A, Deak S B, Riley D J, Boyd C D  
and Stolle C A, (1991) Matrix 11: 359-366.
31. Yeh H, Ornstein-Goldstein N, Indik Z, Sheppard P,  
Anderson N, Rosenbloom J C, Cicila G, Yoon K and Rosenbloom  
J (1987) Collagen Rel. Res. 7: 235-237
- 35 32. Hinek A and Rabinovitch M (1994) J. Cell Biol. 126:



- 563-574.
33. Davidson J M and Sephel G C (1987) *Meth. Enzym.* 144: 214-232.
34. Mecham R P, Hinek A, Entwistle R, Wrenn D S, Griffin G  
5 L and Senior R M (1989) *Biochem.* 28: 3716-3722.
35. Foster, J. A. and Curtiss S. W. (1990) *Am. J. Physio.* 259: L13-L23.
36. Ewart A K, Morris C A, Atkinson D, Jin W, Sternes K, Spallone P, Stock A D, Leppert M and Keating M T (1993)  
10 *Nature Gen.* 5: 11-16.
37. Smith D B and Johnson K S (1988) *Gene* 67: 31-40.
38. Tarlton J F, Vickery C J, Leaper D J and Bailey A J (1997) *B. J. Derm.* 137: 506-516.

0943813 04604

~~67~~ 67  
108

Claims:

1. A method for reducing or eliminating the susceptibility of a tropoelastin to proteolysis comprising mutating a sub-sequence in the tropoelastin so that the susceptibility of the tropoelastin to proteolysis is reduced or eliminated.

2. A method according to claim 1 wherein one sub-sequence is mutated.

3. A method according to claim 1 wherein one amino acid residue in the sub-sequence is mutated.

4. A method according to claim 1 wherein the sub-sequence is capable of being digested by a serine protease.

5. A method according to claim 4 wherein the sub-sequence has an amino acid sequence including the sequence: RAAAG.

6. A method according to claim 5 wherein the sub-sequence is mutated by replacing arginine in the sequence: RAAAG with alanine.

7. A method according to claim 4 wherein the sub-sequence has an amino acid sequence selected from the group of sequences shown in SEQ ID NOS: 17 to 44.

8. A method according to claim 7 wherein the sub-sequence is mutated by replacing arginine in the sequence selected from the group of sequences shown in SEQ ID NOS: 17 to 44 with alanine.

9. A method according to claim 4 wherein the sub-sequence is capable of being digested by thrombin and has an amino acid sequence shown in SEQ ID NOS: 8 or 9.

10. A method according to claim 4 wherein the sub-sequence is capable of being digested by plasmin and has an amino acid sequence shown in SEQ ID NOS: 11 or 12.

11. A method according to claim 4 wherein the sub-sequence is capable of being digested by kallikrein.

12. A method according to claim 11 wherein the sub-sequence has an amino acid sequence shown in SEQ ID NOS: 9 or 10.

13. A method according to claim 1 wherein the sub-  
sequence is capable of being digested by a  
metalloproteinase.

14. A method according to claim 13 wherein the sub-  
5 sequence has an amino acid sequence including the sequence:  
ALAAA.

15. A method according to claim 14 wherein the sub-  
sequence is mutated by replacing alanine at any position in  
the sequence: ALAAA with another amino acid residue.

10 16. A method according to claim 15 wherein the sub-  
sequence is mutated by replacing the alanine which is N-  
terminal to leucine in the sequence: ALAAA with another  
amino acid.

15 17. A method according to claim 13 wherein the sub-  
sequence has an amino acid sequence selected from the group  
of sequences shown in SEQ ID NOS: 45 to 70.

18. A method according to claim 17 wherein the sub-  
sequence is mutated by replacing alanine at any position in  
the sequence selected from the group of sequences shown in  
20 SEQ ID NOS: 45 to 70 with another amino acid residue.

19. A method according to claim 18 wherein the alanine  
that is replaced is N-terminal to leucine.

20. A method according to claim 13 wherein the sub-  
sequence is capable of being digested by gelatinase A or B.

25 21. A method according to claim 20 wherein the sub-  
sequence has an amino acid sequence shown in SEQ ID NO: 13.

22. A method according to any one of claims 1 to 21  
wherein the tropoelastin is human tropoelastin.

30 23. A method for enhancing the susceptibility of a  
tropoelastin to proteolysis comprising inserting a sub-  
sequence into the tropoelastin so that the susceptibility  
of the tropoelastin to proteolysis is enhanced.

24. A method according to claim 23 wherein one sub-  
sequence is inserted.

35 25. A method according to claim 23 wherein the  
inserted sub-sequence is capable of being digested with a

69  
110

serine protease.

26. A method according to claim 25 wherein the inserted sub-sequence has an amino acid sequence including the sequence: RAAAG.

5 27. A method according to claim 25 wherein the inserted sub-sequence has an amino acid sequence selected from the group of sequences shown in SEQ ID NOS: 17 to 44.

28. A method according to claim 25 wherein the inserted sub-sequence is capable of being digested by  
10 thrombin and has an amino acid sequence shown in SEQ ID NOS: 8 or 9.

29. A method according to claim 25 wherein the inserted sub-sequence is capable of being digested by plasmin and has an amino acid sequence shown in SEQ ID NOS:  
15 11 or 12.

30. A method according to claim 25 wherein the inserted sub-sequence is capable of being digested by kallikrein.

31. A method according to claim 30 wherein the  
20 inserted sub-sequence has an amino acid sequence shown in SEQ ID NOS: 9 or 10.

32. A method according to claim 23 wherein the inserted sub-sequence is capable of being digested by a metalloproteinase.

25 33. A method according to claim 32 wherein the inserted sub-sequence has an amino acid sequence including the sequence: ALAAA.

34. A method according to claim 32 wherein the inserted sub-sequence has an amino acid sequence selected  
30 from the group of sequences shown in SEQ ID NOS: 45 to 70.

35. A method according to claim 32 wherein the inserted sub-sequence is capable of being digested by gelatinase A or B.

36. A method according to claim 35 wherein the  
35 inserted sub-sequence has the amino acid sequence shown in SEQ ID NO: 13.

09743818.042601

~~70~~ 70

37. A method according to any one of claims 23 to 36 wherein the tropoelastin is human tropoelastin.

38. A peptidomimetic molecule comprising all or part of a peptide selected from the group consisting of  
5 KAPGVGGAF, RAAAGLG, RLSPELREGD, KAAQFGLVPGV, KSAAKVAAKAQLRAA, RLSPELRE and LAAAKAAKYGAA.

39. A peptidomimetic molecule which has the sequence:  
H-Ala-Ala-Lys-Ala-Gln-Leu-Arg-Ala-Ala-Ala-Gly-Leu-Gly-Ala-  
OH or H-Ala-Ala-Lys-Ala-Gln-Leu-Arg-R-Ala-Ala-Ala-Gly-Leu-  
10 Gly-Ala-OH (where R = a reduced peptide bond).

40. A peptidomimetic molecule which is a retro-inverso pseudo peptide which has the sequence: H-D-Ala-Gly-D-Leu-Gly-D-Ala-D-Ala-D-Ala-(R)-D-Arg-D-Leu-D-Gln-D-Ala-D-Lys-D-Ala-D-Ala-OH (where R = a reduced peptide bond) or H-D-Ala-Gly-D-Leu-Gly-D-Ala-D-Ala-D-Ala-D-Arg-D-Leu-D-Gln-D-Ala-D-Lys-D-Ala-D-Ala-OH.  
15

41. A peptidomimetic molecule which has the sequence H-Val-Pro-Gly-Ala-Leu-Ala-Ala-Ala-OH or H-Val-Pro-Gly-Ala-(R)-Leu-Ala-Ala-Ala-OH (where R = a reduced peptide bond)  
20

42. A peptidomimetic molecule which is a retro-inverso pseudo peptide which has the sequence:: H-D-Ala-D-Ala-D-Ala-D-Leu-(R)-D-Ala-Gly-D-Pro-D-Val-OH (where R = a reduced peptide bond) or H-D-Ala-D-Ala-D-Ala-D-Leu-D-Ala-Gly-D-Pro-D-Val-OH.

43. A method for enhancing the purification of a tropoelastin comprising including a peptidomimetic molecule according to any one of claims 38 to 42 in a crude tropoelastin preparation which is being subjected to purification.  
25

44. A pharmaceutical composition comprising a peptidomimetic molecule according to any one of claims 38 to 42 and a pharmaceutically acceptable carrier.  
30

Abstract

The present invention relates to: manipulation of the amino acid sequence of tropoelastin, particularly human  
5 tropoelastin, to modify its protease susceptibility; to tropoelastin derivatives having modified protease susceptibility; to peptidomimetic molecules which contain amino acid sequences which correspond to or incorporate the protease susceptible sequences of tropoelastin; and to uses  
10 of the tropoelastin derivatives and peptidomimetic molecules.

The invention also relates to nucleic acid molecules and genetic constructs encoding the amino acid sequences of the derivatives and peptidomimetic molecules of the  
15 invention.

09743348.042604




Figure 1

2/26

1 GATCCATGGGTGGCGTTCCGGGTGCTATCCCGGGTGGCGTTCCGGGTGGTGTATTCTACC 60  
GTACCCACCGCAAGGCCACGATAGGGCCACCGCAAGGCCACACATAAGATGG  
S M G G V P G A I P G G V P G G V F Y P  
↑  
Start pf mature processed protein

61 CAGGCGCGGGTCTGGGTGCACTGGGCGGTGGTGGCTGGGCCCGGGTGGTAAACCGCTGA 120  
GTCCGCGCCCAGACCCACGTGACCCGCCACACGCGACCCGGGCCCACCATTGCGCGACT  
G A G L G A L G G G A L G P G G K P L K

121 AACCGGTTCCAGGCGGTCTGGCAGGTGCTGGTCTGGGTGCAAGTCTGGGCGCGTTCCCGG 180  
TTGGCCAAGGTCCGCCAGACCGTCCACGACCAGACCCACGTCCAGACCCGCGCAAGGGCC  
P V P G G L A G A G L G A G L G A F P A

181 CGGTTACCTTCCCGGGTGTCTGGTTCGGGTGGCGTTGCAGACGCAGCTGCTGCGTACA 240  
GCCAATGGAAGGGCCCACGAGACCAAGGGCCACCGCAACGTCTGCGTCGACGACGCATGT  
V T F P G A L V P G G V A D A A A A Y K

241 AAGCGGCAAAGGCAGGTGCGGGTCTGGGCGGGGTACCAGGTGTTGGCGGTCTGGGTGTAT 300  
TTCGCCGTTTCCGTCCACGCCCAGACCCGCCCCATGGTCCACAACCGCCAGACCCACATA  
A A K A G A G L G G V P G V G G L G V S

301 CTGCTGGCGCAGTTGTTCCGACGCCGGGTGCAGGTGTAAAACCGGGCAAAGTTCAGGTG 360  
GACGACCGCGTCAACAAGGCGTCGGGCCACGTCCACATTTGGCCCGTTTCAAGGTCCAC  
A G A V V P Q P G A G V K P G K V P G V

361 TTGGTCTGCCGGGCGTATACCGGGTGGTGTCTGCCGGGCGCGCGTTTCCAGGTGTTG 420  
AACCAGACGGCCGCATATGGGCCCCACCACAAGACGGCCCGCGCGCAAAGGTCCACAAC  
G L P G V Y P G G V L P G A R F P G V G

Figure 2(a)



3/26

421 GTGTACTGCCGGGCGTTCCGACCGGTGCAGGTGTTAAACCGAAGGCACCAGGTGTAGGCG 480  
CACATGACGGCCCGCAAGGCTGGCCACGTCCACAATTTGGCTTCCGTGGTCCACATCCGC  
V L P G V P T G A G V K P K A P G V G G

481 GCGCGTTTCGCGGGTATCCCGGGTGTGGCCCGTTTCGGTGGTCCGCAGCCAGGCGTTCCGC 540  
CGCGCAAGCGCCCATAGGGCCCCACAACCGGGCAAGCCACCAGGCGTCGGTCCGCAAGGCG  
A F A G I P G V G P F G G P Q P G V P L

541 TGGGTTACCCGATCAAAGCGCCGAAGCTTCCAGGTGGCTACGGTCTGCCGTACACCACCG 600  
ACCAATGGGCTAGTTTCGCGGCTTCGAAGGTCCACCGATGCCAGACGGCATGTGGTGGC  
G Y P I K A P K L P G G Y G L P Y T T G

601 GTAAACTGCCGTACGGCTACGGTCCGGGTGGCGTAGCAGGTGCTGCGGGTAAAGCAGGCT 660  
CATTTGACGGCATGCCGATGCCAGGCCACCGCATCGTCCACGACGCCCATTTTCGTCCGA  
K L P Y G Y G P G G V A G A A G K A G Y

661 ACCCAACCGGTACTGGTGTGGTCCGCAGGCTGCTGCGGCAGCTGCGGGCAAGGCAGCAG 720  
TGGGTTGGCCATGACCACAACCGGCTCCGACGACGCCGTGACGCCGCTTCCGTTCGTC  
P T G T G V G P Q A A A A A A A K A A A

721 CAAAATTTCGGCGCGGGTGCAGCGGGTGTCTGCGGGGCGTAGGTGGTGTGGCGTTCCGG 780  
GTTTTAAGCCGCGCCACGTGCGCCACAAGACGGCCCGCATCCACCACGACCGCAAGGCC  
K F G A G A A G V L P G V G G A G V P G

781 GTGTTCCAGGTGCGATCCCGGGCATCGGTGGTATCGCAGGCGTAGGTACTCCGGCGGGCCG 840  
CACAAGGTCCACGCTAGGGCCCGTAGCCACCATAGCGTCCGCATCCATGAGGCCGCCGCG  
V P G A I P G I G G I A G V G T P A A A

841 CTGCGGCTGCGGCAGCTGCGGCGAAAGCAGCTAAATACGGTGGCGGCAGCAGGCCCTGGTTC 900  
GACGCCGACGCCGTGACGCCGCTTTCGTGATTTATGCCACGCCGTCGTCCGGACCAAG  
A A A A A A A K A A K Y G A A A G L V P

Figure 2(b)

901 CGGGTGGTCCAGGCTTCGGTCCGGGTGTTGTAGGCGTTCCGGGTGCTGGTGTTCGGGGCG 960  
GCCACCAGGTCCGAAGCCAGGCCCAACATCCGCAAGGCQACGACCACAAGGCCCGC  
G G P G F G P G V V G V P G A G V P G V

961 TAGGTGTTCCAGGTGCGGGCATCCCGGTTGTACCGGGTGCAGGTATCCCGGGCGCTGCGG 1020  
ATCCACAAGGTCCACGCCCGTAGGGCCAACATGGCCACGTCCATAGGGCCCGCGACGCC  
G V P G A G I P V V P G A G I P G A A V

1021 TTCCAGGTGTTGTATCCCCGGAAGCGGCAGCTAAGGCTGCTGCGAAAGCTGCGAAATACG 1080  
AAGGTCCACACATAGGGGCTTCGCGCGTCGATTCCGACGACGCTTTCGACGCTTTATGC  
P G V V S P E A A A K A A A K A A K Y G

1081 GAGCTCGTCCGGGCGTTGGTGTGGTGGCATCCCGACCTACGGTGTAGGTGCAGGCGGTT 1140  
CTCGAGCAGGCCCGCAACCACAACCACCGTAGGGCTGGATGCCACATCCACGTCCGCCAA  
A R P G V G V G G I P T Y G V G A G G F

1141 TCCAGGTTTCGGCGTTGGTGTGGTGGCATCCCGGGTGTAGCTGGTGTTCGGTCTGTGG 1200  
AGGGTCCAAAGCCGCAACCACAACCACCGTAGGGCCACATCGACCACAAGGCAGACAAC  
P G F G V G V G G I P G V A G V P S V G

1201 GTGGCGTACCGGGTGTGGTGGCGTTCCAGGTGTAGGTATCTCCCCGGAAGCGCAGGCAG 1260  
CACC~~CG~~CATGGGCCACAACCACCGCAAGGTCCACATCCATAGAGGGGCGCTTCGCGTCCGTC  
G V P G V G G V P G V G I S P E A Q A A

1261 CTGCGGCAGCTAAAGCAGCGAAGTACGGCGTTGGTACTCCGGCGGCAGCAGCTGCTAAAG 1320  
GACGCCGTGATTTCGTGCGTTTCATGCCGCAACCATGAGGCCGCGCTCGTCGACGATTTC  
A A A K A A K Y G V G T P A A A A A K A

1321 CAGCGGCTAAAGCAGCGCAGTTCGGACTAGTTCGGGGCGTAGGTGTTGCGCCAGGTGTTG 1380  
GTCGCCGATTTCGTGCGGTCAAGCCTGATCAAGGCCCGCATCCACACGCGGTCCACAAC  
A A K A A Q F G L V P G V G V A P G V G

Figure 2(c)

1381 GCGTAGCACCGGGTGTGGTGTGCTCCGGGCGTAGGTCTGGCACCGGGTGTGGCGTTG 1440  
CGCATCGTGGCCCAACCAACGAGGCCCGCATCCAGACCGTGGCCCAACCGCAAC  
V A P G V G V A P G V G L A P G V G V A

1441 CACCAGGTGTAGGTGTGCGCCGGGCGTTGGTGTAGCACCGGGTATCGGTCCGGGTGGCG 1500  
GTGGTCCACATCCACAACGCGGCCGCAACCACATCGTGGCCCATAGCCAGGCCACCGC  
P G V G V A P G V G V A P G I G P G G V

1501 TTGCGGCTGCTGCGAAATCTGCTGCGAAGGTGCTGCGAAAGCGCAGCTGCGTGCAGCAG 1560  
AACGCCGACGACGCTTTAGACGACGCTTCCAACGACGCTTTCGCGTCGACGCACGTCGTC  
A A A A K S A A K V A A K A Q L R A A A

1561 CTGGTCTGGGTGCGGGCATCCCAGGTCTGGGTGTAGGTGTGGTGTTCGGGGCCTGGGTG 1620  
GACCAGACCCACGCGCGTAGGGTCCAGACCCACATCCACAACCAAGGCCCGGACCCAC  
G L G A G I P G L G V G V G V P G L G V

1621 TAGGTGCAGGGGTACCGGGCCTGGGTGTGGTGCAGGCGTTCCGGGTTTCGGTGTGCGG 1680  
ATCCACGTCCCATGGCCCGGACCCACAACCACGTCCGCAAGGCCCAAAGCCACGACCGC  
G A G V P G L G V G A G V P G F G A G A

1681 CGGACGAAGGTGTACGTGCTCCCTGTCTCCAGAACTGCGTGAAGGTGACCCGTCTCTTT 1740  
GCCTGCTTCCACATGCAGCAAGGGACAGAGGTCTTGACGCACTTCCACTGGGCAGGAGAA  
D E G V R R S L S P E L R E G D P S S S

1741 CCCAGCACCTGCGGTCTACCCCGTCTCTCCAGTGTTCGGGGCGCGCTGGCTGCTGCCA 1800  
GGGTGCTGGACGGCAGATGGGGCAGGAGAGGTGCACAAGGCCCGCGCAGCCGACGACGCT  
Q H L P S T P S S P R V P G A L A A A K

1801 AAGCGGCGAAATACGGTGCAGCGGTTCGGGTGTACTGGGCGGTCTGGGTGCTCTGGGCG 1860  
TTGCGCGCTTTATGCCACGTGCCAAGGCCACATGACCCGCCAGACCCACGAGACCCGC  
A A K Y G A A V P G V L G G L G A L G G

Figure 2(d)

6/26

1861 GTGTTGGTATCCCGGGCGGTGTTGTAGGTGCAGGCCAGCTGCAGCTGCTGCTGCGGCAA 1920  
CACAACCATAGGGCCCGCCACAACATCCAGTCCGGGTCGACGTCGACGACGACGCCGTT  
V G I P G G V V G A G P A A A A A A A K

1921 AGGCAGCGGCGAAAGCAGCTCAGTTCGGTCTGGTTGGTGCAGCAGGTCTGGGCGGTCTGG 1980  
TCCGTCGCGCGCTTTCGTTCGAGTCAAGCCAGACCAACCACGTCGTCAGACCCGCCAGACC  
A A A K A A Q F G L V G A A G L G G L G

1981 GTGTTGGCGGTCTGGGTGTACCGGGCGTTGGTGGTCTGGGTGGCATCCCGCCGGCGGCGG 2040  
CACAACCGCCAGACCCACATGGCCCGCAACCACCAGACCCACCGTAGGGCGGCGCGCGCC  
V G G L G V P G V G G L G G I P P A A A

2041 CAGCTAAAGCGGCTAAATACGGTGCAGCAGGTCTGGGTGGCGTTCTGGGTGGTGTCTGGTC 2100  
GTCGATTTGCGCGATTATGCCACGTCGTCAGACCCACCGCAAGACCCACCACGACCAG  
A K A A K Y G A A G L G G V L G G A G Q

2101 AGTTCCTTGGGCGGTGTAGCGGCACGTCCGGGTTTCGGTCTGTCCCCGATCTTCCAG 2160  
TCAAGGGTGACCCGCCACATCGCCGTGCAGGCCCAAAGCCAGACAGGGGCTAGAAGGGTC  
F P L G G V A A R P G F G L S P I F P G

2161 GCGGTGCATGCCTGGGTAAAGCTTGCGGCCGTAAACGTAAATAATGATAG 2210  
CGCCACGTACGGACCCATTTGAAACGCGGCATTTCGATTTATTACTATCCTAG  
G A C L G K A C G R K R K \* \* \*

Figure 2(e)

7/26

```

1  GGVPGAIPGGVPGGVFFPGAGLGAALGGGALGPGGKPLKPVPGGLAGAGLG 50
  |||
1  GGVPGAIPGGVPGGVFFPGAGLGAALGGGALGPGGKPLKPVPGGLAGAGLG 50
51  AGLGAFPAVTFPGALVPGGVADAAAAYKAAKAGAGLGGVPGVGGGLGVSAG 100
  |||
51  AGLGAFPAVTFPGALVPGGVADAAAAYKAAKAGAGLGGVPGVGGGLGVSAG 100
101  AVVPOPGAGVKPGKVPGVGLPGVYPGGVLPGARFPGVGVLPGVPTGAGVK 150
  |||
101  AVVPOPGAGVKPGKVPGVGLPGVYPGGVLPGARFPGVGVLPGVPTGAGVK 150
151  PKAPGVGGAFAGIPGVGPFGGPQPGVPLGYPIKAPKLPGGYGLPYTTGKL 200
  |||
151  PKAPGVGGAFAGIPGVGPFGGPQPGVPLGYPIKAPKLPGGYGLPYTTGKL 200
201  FYGYGPGGGVAGAAAGKAGYPTGTGVGPQAAAAAATAAAKFGAGAAGVLP 250
  |||
201  FYGYGPGGGVAGAAAGKAGYPTGTGVGPQAAAAAATAAAKFGAGAAGVLP 250
251  VGGAGVPGVPGAIPGIGGIAGVGTAAAAAATAAAKAAKYGAAGGLVPGG 300
  |||
251  VGGAGVPGVPGAIPGIGGIAGVGTAAAAAATAAAKAAKYGAAGGLVPGG 300
301  PGFGPGVVGVPAGVPGVPGAGIPVVPAGIPGAAPVGVVSPEAAAKA 350
  |||
301  PGFGPGVVGVPAGVPGVPGAGIPVVPAGIPGAAPVGVVSPEAAAKA 350
351  AAKAAKYGARPGVGVGGIPTYGVGAGGFPGRGVGVGGIPGVAGVPSVGGV 400
  |||
351  AAKAAKYGARPGVGVGGIPTYGVGAGGFPGRGVGVGGIPGVAGVPSVGGV 400
401  PGVGGVPGVGISPEAQAAAAAATAAAKYGVGTAAAAAATAAAKAAQFGLVPG 450
  |||
401  PGVGGVPGVGISPEAQAAAAAATAAAKYGVGTAAAAAATAAAKAAQFGLVPG 450
451  VGVAPGVGVAPGVGVAPGVGLAPGVGVAPGVGVAPGVGVAPGVGPGGVA 500
  |||
451  VGVAPGVGVAPGVGVAPGVGLAPGVGVAPGVGVAPGVGVAPGVGPGGVA 500
501  AAKSAAKVAAKAAQRAAAGLGAAGIPGLGVGVGVPGGLGVGAGVPGLVGAG 550
  |||
501  AAKSAAKVAAKAAQRAAAGLGAAGIPGLGVGVGVPGGLGVGAGVPGLVGAG 550
551  VPGFGAGADEGVRRSLSPFLREGDPSSSQHLPTSPSPRVPGALAAAKAA 600
  |||
551  VPGFGA.....VPGALAAAKAA 567
601  KYGAAPVGVGLGGLGALGVGIPGGVVGAGEAAAAAATAAAKAAQFGLV 650
  |||
568  KYGAAPVGVGLGGLGALGVGIPGGVVGAGEAAAAAATAAAKAAQFGLV 617
651  AAGLGGGLGVGGGVPGVGGGLGIPAAAAAATAAAKYGAAGLGGVGLGAGOFF 700
  |||
618  AAGLGGGLGVGGGVPGVGGGLGIPAAAAAATAAAKYGAAGLGGVGLGAGOFF 667
701  LGGVAARPGEGLSPITPGGACLGKACGRKK 731
  |||
668  LGGVAARPGEGLSPITPGGACLGKACGRKK 698

```

Figure 3

SUBSTITUTE SHEET (Rule 26) (RO/AU)

09743818.042601

1 ATGGGTGGCGTTCCGGGTGCTGTTCCGGGTGGCGTTCCGGGTGGGTATT 50  
 1 MetGlyGlyValProGlyAlaValProGlyGlyValProGlyGlyValPh 17  
 51 CTACCCAGGCGCGGGTTCCGGTGTGTTCCGGGTGGCGTTCCAGACGAG 100  
 18 eTyrProGlyAlaGlyPheGlyAlaValProGlyGlyValAlaAspAla 34  
 101 CTGCTGCGGTACAAAGCGGCAAAAGCAGGTGCGGGTCTGGGGCGGGTACCA 150  
 35 laAlaAlaTyrLysAlaAlaLysAlaGlyAlaGlyLeuGlyGlyValPro 50  
 151 GGTGTGGCGGTCTGGGTGTATCTGCTGGCGCAGTTGTTCCGCAGCGGG 200  
 51 GlyValGlyGlyLeuGlyValSerAlaGlyAlaValValProGlnProG 67  
 201 TGCAGGTGTAAACCGGGCAAAAGTTCAGGTGTGGTCTGCGGGCGGTAT 250  
 68 yAlaGlyValLysProGlyLysValProGlyValGlyLeuProGlyValT 84  
 251 ACCCGGGTTCCGGTGTGTTCCGGGTGGCGGTTCAGGTGTGGGTAT 300  
 85 yrProGlyPheGlyAlaValProGlyAlaArgPheProGlyValGlyVal 100  
 301 CTGCGGGCGGTTCGCGACCGGTGCGAGGTGTAAACCGAAGGCACCGGTGT 350  
 101 LeuProGlyValProThrGlyAlaGlyValLysProLysAlaProGlyVa 117  
 351 AGCGGGCGCGTTCCGGGTATCCCGGTGTGGCGCGGTTCGGTGGTCCGC 400  
 118 lGlyGlyAlaPheAlaGlyIleProGlyValGlyProPheGlyGlyProG 134  
 401 AGCCAGGCGTTCCGGTGTGGGTATCCCGATCAAAGCGCGAAGCTTCAGGT 450  
 135 lnProGlyValProLeuGlyTyrProIleLysAlaProLysLeuProGly 150  
 451 GGCTACGGTCTGCGGTACACACCGGTAAACTGCGGTACGGGTACGGTCC 500  
 151 GlyTyrGlyLeuProTyrThrThrGlyLysLeuProTyrGlyTyrGlyPr 167  
 501 GGTGGCGGTAGCAGGTGTGCGGGTAAAGCAGGCTACCCAAACCGGTACTG 550  
 168 oGlyGlyValAlaGlyAlaAlaGlyLysAlaGlyTyrProThrGlyThrG 184  
 551 GTGTTGGTCCGCGAGGTGCTGCGGCAGGTGCGGCGAAGGCAGCAGCAAAA 600  
 185 lyValGlyProGlnAlaAlaAlaAlaAlaAlaAlaLysAlaAlaLys 200  
 601 TTCGGCGCGGGTGCAGCGGGTTTCGGTGTGTTCCGGGTGGGTGGGTGC 650  
 201 PheGlyAlaGlyAlaAlaGlyPheGlyAlaValProGlyValGlyGlyAl 217  
 651 TGGCGTTCGGGTGTTCAGGTGCGGTATCCGGGCATCGGTGGTATCGCAG 700  
 218 aGlyValProGlyValProGlyAlaAlaProGlyIleGlyGlyIleAlaG 234  
 701 GGTAGGTATCCGGCGCGCGGTGCGGTGCGGCAGGTGCGGCAGCAAGCA 750  
 235 lyValGlyThrProAlaAlaAlaAlaAlaAlaAlaAlaAlaLysAla 250  
 751 GCTAAATACGGTGCAGCAGGCGGTGCGGTTCGGGTGGGTCCAGGCTTCGG 800  
 251 AlaLysTyrGlyAlaAlaAlaGlyLeuValProGlyGlyProGlyPheG 267  
 801 TCCGGGTGTGTAGCGGTTCGGGTTCGGGTGTGTTCCGGGTGGGTGGGTG 850

Figure 4(a)

SUBSTITUTE SHEET (Rule 26) (RO/AU)

09743818-042601

9/26

268 yProGlyValValGlyValProGlyPheGlyAlaValProGlyValGlyV 284  
851 TTCAGGTGCGGGCATCCGGTGTGTACCGGGTGCAGGTATCCCGGGCGGT 900  
|||||  
285 aLProGlyAlaGlyIleProValValProGlyAlaGlyIleProGlyAla 300  
901 GCGGGTTTCGGTGTGTATCCCGGARGCGGCAGCTAAGGCTGCTGCGAA 950  
|||||  
301 AlaGlyPheGlyAlaValSerProGluAlaAlaAlaAlaAlaAlaAla 317  
951 AGCTGCGAAATACGGAGCTCGTCCGGGGCGTGTGTGTGGTGGCATCCCGA 1000  
|||||  
318 sAlaAlaLysTyrGlyAlaArgProGlyValGlyValGlyGlyIleProT 334  
1001 CCTACGGTGTAGGTGCAGGCGGTTTCCAGGTTTCCGGCGTGTGTGTGGT 1050  
|||||  
335 hrTyrGlyValGlyAlaGlyGlyPheProGlyPheGlyValGlyValGly 350  
1051 GGCATCCCGGGTGTAGCTGGTGTTCGGTCTGTGTGGTGGCGTACCGGGTGT 1100  
|||||  
351 GlyIleProGlyValAlaGlyValProSerValGlyGlyValProGlyVa 367  
1101 TGGTGGCGTTCAGGTGTAGGTATCTCCCGGARGCGGCAGGCTGCGG 1150  
|||||  
368 lGlyGlyValProGlyValGlyIleSerProGluAlaGlnAlaAlaAla 384  
1151 CAGCTAARGCAGCGAAGTACGGCGTGGTACTCCCGCGGCAGCAGCTGCT 1200  
|||||  
385 lAlaAlaLysAlaAlaLysTyrGlyValGlyThrProAlaAlaAlaAla 400  
1201 AAAGCAGCGGCTAARGCAGCGCAGTTCGGACTAGTTCGGCGGTAGGTGT 1250  
|||||  
401 LysAlaAlaAlaLysAlaAlaGlnPheGlyLeuValProGlyValGlyVa 417  
1251 TCGCCAGGTGTGTGGCGTAGCACCGGGTGTGGTGTGTGCTCCGGCGGTAG 1300  
|||||  
418 lAlaProGlyValGlyValAlaProGlyValGlyValAlaProGlyValG 434  
1301 GTCTGGCACCGGGTGTGTGGCGTGCACCAGGTGTAGGTGTGTGGCGCGGC 1350  
|||||  
435 lYLeuAlaProGlyValGlyValAlaProGlyValGlyValAlaProGly 450  
1351 GTTGTGTAGCACCGGGTATCGGTCCGGGTGGCGTGTGGCGCTGCTGCGAA 1400  
|||||  
451 ValGlyValAlaProGlyIleGlyProGlyGlyValAlaAlaAlaAla 467  
1401 ATCTGCTGCGAAGGTGTGCTGCGAARGCGCAGCTGCTGCGCAGCAGCTGCT 1450  
|||||  
468 sSerAlaAlaLysValAlaAlaLysAlaGlnLeuArgAlaAlaAlaGlyL 484  
1451 TGGGTGCGGGCATCCAGGTCTGGGTGTAGGTGTGTGTGTTCGGGGCGTG 1500  
|||||  
485 euGlyAlaGlyIleProGlyLeuGlyValGlyValGlyValProGlyLeu 500  
1501 GGTGTAGGTGCAGGGGTACCGGGCGTGGGTGTGGTGTGAGGCGGTTCGGG 1550  
|||||  
501 GlyValGlyAlaGlyValProGlyLeuGlyValGlyAlaGlyValProG 517  
1551 TTTCGGTGTCTGTTCGGGGCGCTGGCTGCTGCGAARGCGGCGAARTACG 1600  
|||||  
518 yPheGlyAlaValProGlyAlaLeuAlaAlaAlaLysAlaAlaLysTyrG 534  
1601 GTGCTGTTCGGGTGTACTGGCGGTCTGGGTGCTCTGGGCGGTGTGTGGT 1650  
|||||  
535 lYAlaValProGlyValLeuGlyGlyLeuGlyAlaLeuGlyGlyValGly 550  
1651 ATCCCGGGCGGTGTGTAGGTGCAGGGCCCGCTGCAGCTGCTGCTGCGGC 1700  
|||||  
551 IleProGlyGlyValValGlyAlaGlyProAlaAlaAlaAlaAlaAla 567

Figure 4(b)

SUBSTITUTE SHEET (Rule 26) (RO/AU)

09/743818-04260

10/26

1701 AAAGGCAGCGCGCAAGCAGCTCAGTTCCGTCCTGGTTGGGTGCAGCAGGTC 1750  
|||||  
568 aLysAlaAlaAlaLysAlaAlaGlnPheGlyLeuValGlyAlaAlaGlyL 584  
1751 TGGGCGGTCCTGGGTCTTGGCGGTCCTGGGTCTACCGGGCGTTGGTGGTCTG 1800  
|||||  
585 euGlyGlyLeuGlyValGlyGlyLeuGlyValProGlyValGlyGlyLeu 600  
1801 GGTGGCATCCCGCGCGCGCGCGCAGCTAAAGCGGCTAAATACGGTGCAGC 1850  
|||||  
601 GlyGlyIleProProAlaAlaAlaAlaLysAlaAlaLysTyrGlyAlaAl 617  
1851 ASGTCTGGGTGGGCTCTCGGCTGGTGGTGGTCTAGTTCCCACTGGGCGGTG 1900  
|||||  
618 aGlyLeuGlyGlyValLeuGlyGlyAlaGlyGlnPheProLeuGlyGlyV 634  
1901 TAGCGGCACGTCGCGGTTTCGGTCTGTCCCGATCTTCCAGGCGGTGCA 1950  
|||||  
635 aAlaAlaArgProGlyPheGlyLeuSerProIlePheProGlyGlyAla 650  
1951 TGCCTGGGTAAAGCTTGGCGCGCTAAACGTAAA 1983  
|||||  
651 CysLeuGlyLysAlaCysGlyArgLysArgLys 661

Figure 4(c)



11/26




Figure 5

12/26




Figure 6

13/26




Figure 7

14/26




Figure 8

15/26




Figure 9

16/26




Figure 10

17/26




Figure 11

18/26




Figure 12



19/26




Figure 13

20/26




Figure 14




Figure 15

22/26

1 2 3




Figure 16

23/26



Figure 17

24/26




Figure 18




Figure 19

26/26




Figure 20



DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare:

That my residence, post office address and citizenship are as stated below next to my name.

That I verily believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **PROTEASE SUSCEPTIBILITY II** the specification of which (check one)

☐

is attached hereto.

☒

was filed on 19/7/99 as International Appl'n Ser. PCT/AU99/00580 and was amended on 24/7/2000

That I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

That I acknowledge the duty to disclose information known to be material to patentability of this application in accordance with Title 37, Code of Federal Regulations §1.56(a).

That I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate on this invention having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

Priority Claimed

☒☐

Yes No

PCT/AU99/00580 PCT 19/7/99  
(Number) (Country) (Day/Month/Year Filed)

PP4723 Australia 17/7/98  
(Number) (Country) (Day/Month/Year Filed)

☒☐

Yes No

(Number) (Country) (Day/Month/Year Filed)

☐☐

Yes No

That I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

United States Application(s)

(Application Serial No.) (Filing Date) (Status)-(Patented, pending, abandoned)

(Application Serial No.) (Filing Date) (Status)-(Patented, pending, abandoned)

(Application Serial No.) (Filing Date) (Status)-(Patented, pending, abandoned)

That all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

I hereby appoint the following attorneys, with full power of substitution and revocation, to prosecute this application and to transact all business in the United States Patent and Trademark Office connected therewith and request that all correspondence and telephone calls in respect to this application be directed to:

HOWSON AND HOWSON, SPRING HOUSE  
CORPORATE CENTER, BOX 457, SPRING HOUSE, PA 19477, Telephone: (215) 540-9210;  
Facsimile: (215) 540-5818

CATHY A KODROFF

REGISTRATION NO.  
33,980.

Full name of sole or one  
joint inventor:

ANTHONY STEVEN WEISS

Inventor's signature:

*Anthony Steven Weiss*

Date:

March 26, 2001

Residence and Post Office Address:

235 RAINBOW STREET

RANDWICK, NEW SOUTH WALES, 2031

AUSTRALIA

*Any*

Citizenship:

AUSTRALIA

Full name of additional joint  
inventor, if any:

Inventor's signature:

Date:

Residence and Post Office Address:

09745918-042601  
T09240-8180

Citizenship: . . . .

Australia

Address for Correspondence:

HOWSON AND HOWSON  
SPRING HOUSE CORPORATE CENTER  
BOX 457  
SPRING HOUSE, PA 19477  
USA

09743813.042601

SEQUENCE LISTING

<110> The University of Sydney

<120> Protease Susceptibility

<130> Weiss Protease

<140>

<141>

<160> 74

<170> PatentIn Ver. 2.0

<210> 1

<211> 2106

<212> DNA

<213> Homo sapiens

<400> 1

atgggtggcg ttccgggtgc tatcccgggt ggcggtccgg gtggtgtatt ctaccaggc 60  
gcgggtcttg gtgactggg cgggtggtgcg ctgggcccgg gtggtaaacc gctgaaaccg 120  
gttccaggcg gtctggcagg tgctggtctg ggtgcaggtc tgggcgcggt cccggcgggt 180  
accttcccgg gtgctctggt tccgggtggc gttgcagacg cagctgctgc gtacaaagcg 240  
gcaaaggcag gtgcgggtct gggcggggta ccagggtgtg gcggtctggg tgtatctgct 300  
ggcgagttg ttccgcagcc ggggtgcagg gtaaaaccgg gcaaagttcc aggtgttggt 360  
ctgccggggc tatacccggg tgggtgttctg ccgggcgcgc gtttcccagg tgttggtgta 420  
ctgccggggc ttccgaccg tgcagggtgt aaaccgaagg caccaggtgt aggcggcgcg 480

09/743818-042601

ttccgcggtta tcccgggtgt tggcccgttc ggtggtccgc agccaggcgt tccgctgggt 540  
 taccgatca aagcgccgaa gcttccaggt ggctacggtc tgccgtacac caccggtaaa 600  
 ctgccgtacg gctacgggtcc ggggtggcgta gcaggtgctg cgggtaaagc aggctaccca 660  
 accggtagctg gtgttggtcc gcaggctgct gcggcagctg cggcgaaggc agcagcaaaa 720  
 ttccgcgcggt gtgcagcggg tgttctgccc ggcgtagggt gtgctggcgt tccgggtgtt 780  
 ccagggtgca tcccgggcat cgggtggatc gcaggcgtag gtactccggc ggccgctgcg 840  
 gctgcggcag ctgcggcgaa agcagctaaa tacgggtgcg cagcaggcct ggttccgggt 900  
 ggtccaggct tcggtccggg tgttgtaggc gttccgggtg ctggtgttcc gggcgtagggt 960  
 gttccagggtg cgggcatccc ggttgtaggc ggtgcaggta tcccgggcgc tgcggttcca 1020  
 ggtgtgttat ccccggaagc ggcagctaag gctgctgca aagctgcgaa atacggagct 1080  
 cgtccggggtg ttggtgttgg tggcatcccg acctacgggt taggtgcagg cggtttccca 1140  
 ggttccggcg ttggtgttgg tggcatcccg ggtgtagctg gtgttccgtc tgttggtggc 1200  
 gtaccgggtg ttggtggcgt tccagggtga ggtatctccc cggaagcgca ggcagctgcg 1260  
 gcagctaaaag cagcgaagta cggcgttggg actccggcgg cagcagctgc taaagcagcg 1320  
 gctaaagcag cgcagttcgg actagttccg ggcgtagggt ttgcgccagg tgttggtgta 1380  
 gcaccgggtg ttggtgttgc tccgggcgta ggtctggcac cgggtgttgg cgttgcacca 1440  
 ggtgtaggtg ttgcgccggg cgttggtgta gcaccgggtg tcgggtccggg tggcgttgcg 1500  
 gctgctgca aatctgctgc gaagggtgct gcgaaagcgc agctgcgtgc agcagctggg 1560  
 ctgggtgctg gcatcccagg tctgggtgta ggtgttgggt ttcgggctt ggggtgtaggt 1620  
 gcaggggtac cgggcctggg tgttggtgca ggcgttccgg gtttcgggtg tgttccgggc 1680  
 gcgctggctg ctgcgaaagc ggcgaaatac ggtgcagcgg ttcgggtgt actgggcggg 1740  
 ctgggtgctc tgggcgggtg tggatccccg ggcgggtgtg taggtgcagg cccagctgca 1800  
 gctgctgctg cggcaaaggc agcggcgaaa gcagctcagt tcgggtcgtt tgggtgcagca 1860  
 ggtctggggtg gtctgggtgt tggcgggtct ggtgtaccgg gcgttggtgg tctgggtggc 1920  
 atcccgccgg cggcggcagc taaagcggct aaatacgggt cagcaggtct ggggtggcgtt 1980  
 ctgggtggtg ctgggtcagtt cccactgggc ggtgtagcgg cacgtccggg tttcgggtctg 2040  
 tccccgatct tcccaggcgg tgcgtgcctg ggtaaagctt gcggccgtaa acgtaaataa 2100  
 tgatag 2106

<210> 2

<211> 1992

<212> DNA

<213> Homo sapiens

<400> 2

atgggtggcg ttccgggtgc tgttccgggt ggcgttccgg gtggtgtatt ctaccaggc 60  
gcgggtttcg gtgctgttcc gggtggcgtt gcagacgcag ctgctgcgta caaagcggca 120  
aaggcagggtg cgggtctggg cggggtacca ggtgttggcg gtctgggtgt atctgctggc 180  
gcagttgttc cgcagccggg tgcagggtga aaaccgggca aagttccagg tgttgggtctg 240  
ccgggcgtat acccgggttt cgggtgctgtt ccgggcgcgc gtttcccagg tgttgggtgta 300  
ctgccgggcg ttccgaccgg tgcagggtgt aaaccgaagg caccagggtgt aggcggcgcg 360  
ttcgcgggta tcccgggtgt tggcccgttc ggtgggtccgc agccaggcgt tccgctgggt 420  
taccgatca aagcgccgaa gcttccagggt ggctacggtc tgccgtacac caccggtaaa 480  
ctgccgtacg gctacgggtcc gggtggcgta gcagggtctg cgggtaaagc aggctacca 540  
accgggtactg gtgttgggtcc gcaggctgct gcggcagctg cggcgaaggc agcagcaaaa 600  
ttcggcgcgg gtgcagcggg ttctgggtgt gttccgggcg taggtggtgc tggcgttccg 660  
ggtgttccag gtgcgatccc gggcatcgggt ggtatcgcag gcgtaggtac tccggcggcc 720  
gctgcggctg cggcagctgc ggcgaaagca gctaaatacg gtgcggcagc aggcctggtt 780  
ccgggtggtc caggcttcgg tccgggtgtt gtaggcgttc cgggtttcgg tgctgttccg 840  
ggcgtaggtg ttccagggtc gggcatcccg gttgtaccgg gtgcagggtat cccgggcgct 900  
gcgggtttcg gtgctgtatc cccggaagcg gcagctaagg ctgctgcgaa agctgcgaaa 960  
tacggagctc gtccgggcgt tgggtgttgggt ggcatcccg cctacgggtgt aggtgcaggc 1020  
ggtttcccag gtttcggcgt tgggtgttgggt ggcatcccg gtgtagctgg tgttccgtct 1080  
gttgggtggcg taccgggtgt tggtggcgtt ccagggtgtag gtatctcccc ggaagcgcag 1140  
gcagctgcgg cagctaaaagc agcgaagtac ggcgttggtgta ctccggcggc agcagctgct 1200  
aaagcagcgg ctaaagcagc gcagttcggga ctagttccgg gcgtaggtgt tgcgccagggt 1260  
gttggcgtag caccgggtgt tgggtgttgt ccgggcgtag gtctggcacc ggggtgttggc 1320

gttgaccag gtgtaggtgt tgcgccgggc gttggtgtag caccgggtat cgggccgggt 1380  
 ggcgttgccg ctgctgcgaa atctgctgcg aaggttgctg cgaaagcgca gctgcgtgca 1440  
 gcagctggtc tgggtgcggg catcccaggt ctgggtgtag gtgttggtgt tccgggcctg 1500  
 ggtgtaggtg caggggtacc gggcctgggt gttggtgcag gcgttcggg tttcgggtgct 1560  
 gttccgggcg cgctggctgc tgcgaaagcg gcgaaatacg gtgctgttcc ggggtgtactg 1620  
 ggcggtcttg gtgctctggg cgggtgttgg atccccggcg gtgttgtagg tgcaggccca 1680  
 gctgcagctg ctgctgcggc aaaggcagcg gcgaaagcag ctgagttcgg tctgggttgg 1740  
 gcagcaggtc tgggcgggtc ggggtgttgg ggtctgggtg taccgggcgt tgggtgtctg 1800  
 ggtggcatcc cgccggcggc ggcagctaaa gcggctaaat acggtgcagc aggtctgggt 1860  
 ggcgttcttg gtggtgctgg tcagttccca ctgggcggtg tagcggcacg tccgggtttc 1920  
 ggtctgtccc cgatcttccc aggcggtgca tgctgggta aagcttgccg ccgtaaacgt 1980  
 aaataatgat ag 1992

<210> 3

<211> 2205

<212> DNA

<213> Homo sapiens

<400> 3

atgggtggcg ttccgggtgc tatccccgggt ggcgttcgg gtggtgtatt ctaccaggc 60  
 gcgggtcttg gtgactggg cgggtgtgcg ctgggcccg gtggtaaacc gctgaaaccg 120  
 gttccaggcg gtctggcagg tgctggtctg ggtgcaggtc tgggcgcgtt cccggcggtt 180  
 accttcccgg gtgctctggt tccgggtggc gttgcagacg cagctgctgc gtacaaagcg 240  
 gcaaaggcag gtgcgggtct gggcggggta ccaggtgttg gcggtctggg tgtatctgct 300  
 ggcgcagttg ttccgcagcc ggggtgcaggt gtaaaaccgg gcaaagttcc aggtgttgg 360  
 ctgccgggcg tataccccgg tggtgttctg ccgggcgcgc gtttcccagg tgttggtgta 420  
 ctgccgggcg ttccgaccgg tgcaggtgtt aaaccgaagg caccaggtgt aggcggcgcg 480  
 ttcgcgggta tcccggtgt tggcccggtc ggtggtccgc agccaggcgt tccgctgggt 540

tacccgatca aagcgccgaa gcttccaggt ggctacggtc tgccgtacac caccggtaaa 600  
ctgccgtacg gctacgggtcc ggggtggcgta gcaggtgctg cgggtaaagc aggctaccca 660  
accgggtactg gtgtttggtcc gcaggctgct gcggcagctg cggcgaaggc agcagcaaaa 720  
ttcggcgcggt gtgcagcggg tgttctgccg ggcgtagggt gtgctggcgt tccgggtggt 780  
ccaggtgcga tcccgggcat cgggtgtatc gcaggcgtag gtactccggc ggccgctgcg 840  
gctgcggcag ctgcggcgaa agcagctaaa tacgggtgcgg cagcaggcct ggttccgggt 900  
ggtccaggct tcgggtccggg tgttgtaggc gttccgggtg ctggtgttcc gggcgtagggt 960  
gttccagggt cgggcatccc ggttgtaggc ggtgcaggta tcccgggcgc tgcggttcca 1020  
ggtgttgtagt ccccggaagc ggcagctaac gctgctgcga aagctgcgaa atacggagct 1080  
cgtccggggtg ttggtgttgg tggcatcccg acctacgggt taggtgcagg cggtttccca 1140  
ggtttcggcg ttggtgttgg tggcatcccg ggtgtagctg gtgttccgct tgttggtggc 1200  
gtaccgggtg ttggtggcgt tccagggtgta ggtatctccc cggaagcgca ggcagctgcg 1260  
gcagctaaag cagcgaagta cggcgttggg actccggcgg cagcagctgc taaagcagcg 1320  
gctaaagcag cgcagttcgg actagttccg ggcgtagggt ttgcgccagg tgttggtgta 1380  
gcaccgggtg ttggtgttgc tccgggcgta ggtctggcac cgggtgttgg cgttgcacca 1440  
ggtgtaggtg ttgcgccggg cgttggtgta gcaccgggtg tcgggtccggg tggcgttgcg 1500  
gctgctgcga aatctgctgc gaaggttgcg gcgaaagcgc agctgcgtgc agcagctggt 1560  
ctgggtgcgg gcatcccagg tctgggtgta ggtgttgggt ttccgggcct ggggtgtaggt 1620  
gcaggggtac cgggcctggg tgttggtgca ggcgttccgg gtttcgggtg tggcgcgga 1680  
gaaggtgtac gtcgttccct gtctccagaa ctgcgtgaag gtgaccgctc ctcttcccag 1740  
cacctgccgt ctaccccgct ctctccaggt gttccggggt cgctgggtgc tgcgaaagcg 1800  
gcgaaatacg gtgcagcggg tccgggtgta ctgggcgggt tgggtgctct gggcgggtgtt 1860  
ggtatcccgg gcggtgttgt aggtgcaggc ccagctgcag ctgctgctgc ggcaaaggca 1920  
gcggcgaaaag cagctcagtt cgggtctggtt ggtgcagcag gtctgggcgg tctgggtgtt 1980  
ggcgggtctg gtgtaccggg cgttggtggt ctgggtggca tcccgcgggc ggcggcagct 2040  
aaagcggcta aatacgggtc agcaggtctg ggtggcggtc tgggtggtgc tggtcagttc 2100  
ccactggggt gtgtagcggt acgtccgggt ttcgggtctgt ccccgatctt cccaggcgggt 2160  
gcatgcctgg gtaaagcttg cggccgtaaa cgtaaataat gatag 2205



<210> 4

<211> 731

<212> PRT

<213> Homo sapiens

<400> 4

Gly Gly Val Pro Gly Ala Ile Pro Gly Gly Val Pro Gly Gly Val Phe

1

5

10

15

Tyr Pro Gly Ala Gly Leu Gly Ala Leu Gly Gly Gly Ala Leu Gly Pro

20

25

30

Gly Gly Lys Pro Leu Lys Pro Val Pro Gly Gly Leu Ala Gly Ala Gly

35

40

45

Leu Gly Ala Gly Leu Gly Ala Phe Pro Ala Val Thr Phe Pro Gly Ala

50

55

60

Leu Val Pro Gly Gly Val Ala Asp Ala Ala Ala Tyr Lys Ala Ala

65

70

75

80

Lys Ala Gly Ala Gly Leu Gly Gly Val Pro Gly Val Gly Gly Leu Gly

85

90

95

Val Ser Ala Gly Ala Val Val Pro Gln Pro Gly Ala Gly Val Lys Pro

100

105

110

Gly Lys Val Pro Gly Val Gly Leu Pro Gly Val Tyr Pro Gly Gly Val

09243318 042601

115

120

125

Leu Pro Gly Ala Arg Phe Pro Gly Val Gly Val Leu Pro Gly Val Pro

130

135

140

Thr Gly Ala Gly Val Lys Pro Lys Ala Pro Gly Val Gly Gly Ala Phe

145

150

155

160

Ala Gly Ile Pro Gly Val Gly Pro Phe Gly Gly Pro Gln Pro Gly Val

165

170

175

Pro Leu Gly Tyr Pro Ile Lys Ala Pro Lys Leu Pro Gly Gly Tyr Gly

180

185

190

Leu Pro Tyr Thr Thr Gly Lys Leu Pro Tyr Gly Tyr Gly Pro Gly Gly

195

200

205

Val Ala Gly Ala Ala Gly Lys Ala Gly Tyr Pro Thr Gly Thr Gly Val

210

215

220

Gly Pro Gln Ala Ala Ala Ala Ala Ala Lys Ala Ala Ala Lys Phe

225

230

235

240

Gly Ala Gly Ala Ala Gly Val Leu Pro Gly Val Gly Gly Ala Gly Val

245

250

255

Pro Gly Val Pro Gly Ala Ile Pro Gly Ile Gly Gly Ile Ala Gly Val

260

265

270

[illegible]

Gly Thr Pro Ala Ala Ala Ala Ala Ala Ala Ala Ala Lys Ala Ala  
275 280 285

Lys Tyr Gly Ala Ala Ala Gly Leu Val Pro Gly Gly Pro Gly Phe Gly  
290 295 300

Pro Gly Val Val Gly Val Pro Gly Ala Gly Val Pro Gly Val Gly Val  
305 310 315 320

Pro Gly Ala Gly Ile Pro Val Val Pro Gly Ala Gly Ile Pro Gly Ala  
325 330 335

Ala Val Pro Gly Val Val Ser Pro Glu Ala Ala Ala Lys Ala Ala Ala  
340 345 350

Lys Ala Ala Lys Tyr Gly Ala Arg Pro Gly Val Gly Val Gly Gly Ile  
355 360 365

Pro Thr Tyr Gly Val Gly Ala Gly Gly Phe Pro Gly Phe Gly Val Gly  
370 375 380

Val Gly Gly Ile Pro Gly Val Ala Gly Val Pro Ser Val Gly Gly Val  
385 390 395 400

Pro Gly Val Gly Gly Val Pro Gly Val Gly Ile Ser Pro Glu Ala Gln  
405 410 415

Ala Ala Ala Ala Ala Lys Ala Ala Lys Tyr Gly Val Gly Thr Pro Ala

09743338 042501

0943318 042601  
10240 "BFB" 420

420

425

430

Ala Ala Ala Ala Lys Ala Ala Ala Lys Ala Ala Gln Phe Gly Leu Val

435

440

445

Pro Gly Val Gly Val Ala Pro Gly Val Gly Val Ala Pro Gly Val Gly

450

455

460

Val Ala Pro Gly Val Gly Leu Ala Pro Gly Val Gly Val Ala Pro Gly

465

470

475

480

Val Gly Val Ala Pro Gly Val Gly Val Ala Pro Gly Ile Gly Pro Gly

485

490

495

Gly Val Ala Ala Ala Ala Lys Ser Ala Ala Lys Val Ala Ala Lys Ala

500

505

510

Gln Leu Arg Ala Ala Ala Gly Leu Gly Ala Gly Ile Pro Gly Leu Gly

515

520

525

Val Gly Val Gly Val Pro Gly Leu Gly Val Gly Ala Gly Val Pro Gly

530

535

540

Leu Gly Val Gly Ala Gly Val Pro Gly Phe Gly Ala Gly Ala Asp Glu

545

550

555

560

Gly Val Arg Arg Ser Leu Ser Pro Glu Leu Arg Glu Gly Asp Pro Ser

565

570

575

Ser Ser Gln His Leu Pro Ser Thr Pro Ser Ser Pro Arg Val Pro Gly  
580 585 590

Ala Leu Ala Ala Ala Lys Ala Ala Lys Tyr Gly Ala Ala Val Pro Gly  
595 600 605

Val Leu Gly Gly Leu Gly Ala Leu Gly Gly Val Gly Ile Pro Gly Gly  
610 615 620

Val Val Gly Ala Gly Pro Ala Ala Ala Ala Ala Ala Ala Lys Ala Ala  
625 630 635 640

Ala Lys Ala Ala Gln Phe Gly Leu Val Gly Ala Ala Gly Leu Gly Gly  
645 650 655

Leu Gly Val Gly Gly Leu Gly Val Pro Gly Val Gly Gly Leu Gly Gly  
660 665 670

Ile Pro Pro Ala Ala Ala Ala Lys Ala Ala Lys Tyr Gly Ala Ala Gly  
675 680 685

Leu Gly Gly Val Leu Gly Gly Ala Gly Gln Phe Pro Leu Gly Gly Val  
690 695 700

Ala Ala Arg Pro Gly Phe Gly Leu Ser Pro Ile Phe Pro Gly Gly Ala  
705 710 715 720

Cys Leu Gly Lys Ala Cys Gly Arg Lys Arg Lys

09243818 042601

725

730

&lt;210&gt; 5

&lt;211&gt; 698

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

Gly Gly Val Pro Gly Ala Ile Pro Gly Gly Val Pro Gly Gly Val Phe

1

5

10

15

Tyr Pro Gly Ala Gly Leu Gly Ala Leu Gly Gly Gly Ala Leu Gly Pro

20

25

30

Gly Gly Lys Pro Leu Lys Pro Val Pro Gly Gly Leu Ala Gly Ala Gly

35

40

45

Leu Gly Ala Gly Leu Gly Ala Phe Pro Ala Val Thr Phe Pro Gly Ala

50

55

60

Leu Val Pro Gly Gly Val Ala Asp Ala Ala Ala Tyr Lys Ala Ala

65

70

75

80

Lys Ala Gly Ala Gly Leu Gly Gly Val Pro Gly Val Gly Gly Leu Gly

85

90

95

Val Ser Ala Gly Ala Val Val Pro Gln Pro Gly Ala Gly Val Lys Pro

100

105

110

09743818-042601  
T09240"8T8E4.60

Gly Lys Val Pro Gly Val Gly Leu Pro Gly Val Tyr Pro Gly Gly Val  
115 120 125

Leu Pro Gly Ala Arg Phe Pro Gly Val Gly Val Leu Pro Gly Val Pro  
130 135 140

Thr Gly Ala Gly Val Lys Pro Lys Ala Pro Gly Val Gly Gly Ala Phe  
145 150 155 160

Ala Gly Ile Pro Gly Val Gly Pro Phe Gly Gly Pro Gln Pro Gly Val  
165 170 175

Pro Leu Gly Tyr Pro Ile Lys Ala Pro Lys Leu Pro Gly Gly Tyr Gly  
180 185 190

Leu Pro Tyr Thr Thr Gly Lys Leu Pro Tyr Gly Tyr Gly Pro Gly Gly  
195 200 205

Val Ala Gly Ala Ala Gly Lys Ala Gly Tyr Pro Thr Gly Thr Gly Val  
210 215 220

Gly Pro Gln Ala Ala Ala Ala Ala Ala Lys Ala Ala Ala Lys Phe  
225 230 235 240

Gly Ala Gly Ala Ala Gly Val Leu Pro Gly Val Gly Gly Ala Gly Val  
245 250 255

092443010374650

Pro Gly Val Pro Gly Ala Ile Pro Gly Ile Gly Gly Ile Ala Gly Val

260

265

270

Gly Thr Pro Ala Ala Ala Ala Ala Ala Ala Ala Ala Lys Ala Ala

275

280

285

Lys Tyr Gly Ala Ala Ala Gly Leu Val Pro Gly Gly Pro Gly Phe Gly

290

295

300

Pro Gly Val Val Gly Val Pro Gly Ala Gly Val Pro Gly Val Gly Val

305

310

315

320

Pro Gly Ala Gly Ile Pro Val Val Pro Gly Ala Gly Ile Pro Gly Ala

325

330

335

Ala Val Pro Gly Val Val Ser Pro Glu Ala Ala Ala Lys Ala Ala Ala

340

345

350

Lys Ala Ala Lys Tyr Gly Ala Arg Pro Gly Val Gly Val Gly Gly Ile

355

360

365

Pro Thr Tyr Gly Val Gly Ala Gly Gly Phe Pro Gly Phe Gly Val Gly

370

375

380

Val Gly Gly Ile Pro Gly Val Ala Gly Val Pro Ser Val Gly Gly Val

385

390

395

400

Pro Gly Val Gly Gly Val Pro Gly Val Gly Ile Ser Pro Glu Ala Gln

405

410

415

0943313.042601  
T09240"STB4260



Ala Ala Ala Ala Ala Lys Ala Ala Lys Tyr Gly Val Gly Thr Pro Ala

420

425

430

Ala Ala Ala Ala Lys Ala Ala Ala Lys Ala Ala Gln Phe Gly Leu Val

435

440

445

Pro Gly Val Gly Val Ala Pro Gly Val Gly Val Ala Pro Gly Val Gly

450

455

460

Val Ala Pro Gly Val Gly Leu Ala Pro Gly Val Gly Val Ala Pro Gly

465

470

475

480

Val Gly Val Ala Pro Gly Val Gly Val Ala Pro Gly Ile Gly Pro Gly

485

490

495

Gly Val Ala Ala Ala Ala Lys Ser Ala Ala Lys Val Ala Ala Lys Ala

500

505

510

Gln Leu Arg Ala Ala Ala Gly Leu Gly Ala Gly Ile Pro Gly Leu Gly

515

520

525

Val Gly Val Gly Val Pro Gly Leu Gly Val Gly Ala Gly Val Pro Gly

530

535

540

Leu Gly Val Gly Ala Gly Val Pro Gly Phe Gly Ala Val Pro Gly Ala

545

550

555

560

094338 04501

Leu Ala Ala Ala Lys Ala Ala Lys Tyr Gly Ala Ala Val Pro Gly Val  
565 570 575

Leu Gly Gly Leu Gly Ala Leu Gly Gly Val Gly Ile Pro Gly Gly Val  
580 585 590

Val Gly Ala Gly Pro Ala Ala Ala Ala Ala Ala Lys Ala Ala Ala  
595 600 605

Lys Ala Ala Gln Phe Gly Leu Val Gly Ala Ala Gly Leu Gly Gly Leu  
610 615 620

Gly Val Gly Gly Leu Gly Val Pro Gly Val Gly Gly Leu Gly Gly Ile  
625 630 635 640

Pro Pro Ala Ala Ala Ala Lys Ala Ala Lys Tyr Gly Ala Ala Gly Leu  
645 650 655

Gly Gly Val Leu Gly Gly Ala Gly Gln Phe Pro Leu Gly Gly Val Ala  
660 665 670

Ala Arg Pro Gly Phe Gly Leu Ser Pro Ile Phe Pro Gly Gly Ala Cys  
675 680 685

Leu Gly Lys Ala Cys Gly Arg Lys Arg Lys  
690 695

<210> 6

09743313 042601  
"09240" BT02460

<211> 660

<212> PRT

<213> Homo sapiens

<400> 6

Gly Gly Val Pro Gly Ala Val Pro Gly Gly Val Pro Gly Gly Val Phe

1 5 10 15

Tyr Pro Gly Ala Gly Phe Gly Ala Val Pro Gly Gly Val Ala Asp Ala

20 25 30

Ala Ala Ala Tyr Lys Ala Ala Lys Ala Gly Ala Gly Leu Gly Gly Val

35 40 45

Pro Gly Val Gly Gly Leu Gly Val Ser Ala Gly Ala Val Val Pro Gln

50 55 60

Pro Gly Ala Gly Val Lys Pro Gly Lys Val Pro Gly Val Gly Leu Pro

65 70 75 80

Gly Val Tyr Pro Gly Phe Gly Ala Val Pro Gly Ala Arg Phe Pro Gly

85 90 95

Val Gly Val Leu Pro Gly Val Pro Thr Gly Ala Gly Val Lys Pro Lys

100 105 110

Ala Pro Gly Val Gly Gly Ala Phe Ala Gly Ile Pro Gly Val Gly Pro

115 120 125

0574336-042604

Phe Gly Gly Pro Gln Pro Gly Val Pro Leu Gly Tyr Pro Ile Lys Ala  
130 135 140

Pro Lys Leu Pro Gly Gly Tyr Gly Leu Pro Tyr Thr Thr Gly Lys Leu  
145 150 155 160

Pro Tyr Gly Tyr Gly Pro Gly Gly Val Ala Gly Ala Ala Gly Lys Ala  
165 170 175

Gly Tyr Pro Thr Gly Thr Gly Val Gly Pro Gln Ala Ala Ala Ala Ala  
180 185 190

Ala Ala Lys Ala Ala Ala Lys Phe Gly Ala Gly Ala Ala Gly Phe Gly  
195 200 205

Ala Val Pro Gly Val Gly Gly Ala Gly Val Pro Gly Val Pro Gly Ala  
210 215 220

Ile Pro Gly Ile Gly Gly Ile Ala Gly Val Gly Thr Pro Ala Ala Ala  
225 230 235 240

Ala Ala Ala Ala Ala Ala Ala Lys Ala Ala Lys Tyr Gly Ala Ala Ala  
245 250 255

Gly Leu Val Pro Gly Gly Pro Gly Phe Gly Pro Gly Val Val Gly Val  
260 265 270

Pro Gly Phe Gly Ala Val Pro Gly Val Gly Val Pro Gly Ala Gly Ile

0943318 042601

275

280

285

Pro Val Val Pro Gly Ala Gly Ile Pro Gly Ala Ala Gly Phe Gly Ala

290

295

300

Val Ser Pro Glu Ala Ala Ala Lys Ala Ala Ala Lys Ala Ala Lys Tyr

305

310

315

320

Gly Ala Arg Pro Gly Val Gly Val Gly Gly Ile Pro Thr Tyr Gly Val

325

330

335

Gly Ala Gly Gly Phe Pro Gly Phe Gly Val Gly Val Gly Gly Ile Pro

340

345

350

Gly Val Ala Gly Val Pro Ser Val Gly Gly Val Pro Gly Val Gly Gly

355

360

365

Val Pro Gly Val Gly Ile Ser Pro Glu Ala Gln Ala Ala Ala Ala Ala

370

375

380

Lys Ala Ala Lys Tyr Gly Val Gly Thr Pro Ala Ala Ala Ala Ala Lys

385

390

395

400

Ala Ala Ala Lys Ala Ala Gln Phe Gly Leu Val Pro Gly Val Gly Val

405

410

415

Ala Pro Gly Val Gly Val Ala Pro Gly Val Gly Val Ala Pro Gly Val

420

425

430

0943838-042601



580

585

590

Val Pro Gly Val Gly Gly Leu Gly Gly Ile Pro Pro Ala Ala Ala Ala

595

600

605

Lys Ala Ala Lys Tyr Gly Ala Ala Gly Leu Gly Gly Val Leu Gly Gly

610

615

620

Ala Gly Gln Phe Pro Leu Gly Gly Val Ala Ala Arg Pro Gly Phe Gly

625

630

635

640

Leu Ser Pro Ile Phe Pro Gly Gly Ala Cys Leu Gly Lys Ala Cys Gly

645

650

655

Arg Lys Arg Lys

660

&lt;210&gt; 7

&lt;211&gt; 571

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 7

Gly Gly Val Pro Gly Ala Ile Pro Gly Gly Val Pro Gly Gly Val Phe

1

5

10

15

Tyr Pro Gly Ala Gly Leu Gly Ala Leu Gly Gly Gly Ala Leu Gly Pro

0943838-042601

20

25

30

Gly Gly Lys Pro Leu Lys Pro Val Pro Gly Gly Leu Ala Gly Ala Gly

35

40

45

Leu Gly Ala Gly Leu Gly Ala Phe Pro Ala Val Thr Phe Pro Gly Ala

50

55

60

Leu Val Pro Gly Gly Val Ala Asp Ala Ala Ala Tyr Lys Ala Ala

65

70

75

80

Lys Ala Gly Ala Gly Leu Gly Gly Val Pro Gly Val Gly Gly Leu Gly

85

90

95

Val Ser Ala Gly Ala Val Val Pro Gln Pro Gly Ala Gly Val Lys Pro

100

105

110

Gly Lys Val Pro Gly Val Gly Leu Pro Gly Val Tyr Pro Gly Gly Val

115

120

125

Leu Pro Gly Ala Arg Phe Pro Gly Val Gly Val Leu Pro Gly Val Pro

130

135

140

Thr Gly Ala Gly Val Lys Pro Lys Ala Pro Gly Val Gly Gly Ala Phe

145

150

155

160

Ala Gly Ile Pro Gly Val Gly Pro Phe Gly Gly Pro Gln Pro Gly Val

165

170

175

T09240"8T884260



Pro Leu Gly Tyr Pro Ile Lys Ala Pro Lys Leu Pro Gly Gly Tyr Gly  
180 185 190

Leu Pro Tyr Thr Thr Gly Lys Leu Pro Tyr Gly Tyr Gly Pro Gly Gly  
195 200 205

Val Ala Gly Ala Ala Gly Lys Ala Gly Tyr Pro Thr Gly Thr Gly Val  
210 215 220

Gly Pro Gln Ala Ala Ala Ala Ala Ala Lys Ala Ala Ala Lys Phe  
225 230 235 240

Gly Ala Gly Ala Ala Gly Val Leu Pro Gly Val Gly Gly Ala Gly Val  
245 250 255

Pro Gly Val Pro Gly Ala Ile Pro Gly Ile Gly Gly Ile Ala Gly Val  
260 265 270

Gly Thr Pro Ala Ala Ala Ala Ala Ala Ala Ala Lys Ala Ala  
275 280 285

Lys Tyr Gly Ala Ala Ala Gly Leu Val Pro Gly Gly Pro Gly Phe Gly  
290 295 300

Pro Gly Val Val Gly Val Pro Gly Ala Gly Val Pro Gly Val Gly Val  
305 310 315 320

Pro Gly Ala Gly Ile Pro Val Val Pro Gly Ala Gly Ile Pro Gly Ala

094333043601

325

330

335

Ala Val Pro Gly Val Val Ser Pro Glu Ala Ala Ala Lys Ala Ala Ala

340

345

350

Lys Ala Ala Lys Tyr Gly Ala Arg Pro Gly Val Gly Val Gly Gly Ile

355

360

365

Pro Thr Tyr Gly Val Gly Ala Gly Gly Phe Pro Gly Phe Gly Val Gly

370

375

380

Val Gly Gly Ile Pro Gly Val Ala Gly Val Pro Ser Val Gly Gly Val

385

390

395

400

Pro Gly Val Gly Gly Val Pro Gly Val Gly Ile Ser Pro Glu Ala Gln

405

410

415

Ala Ala Ala Ala Ala Lys Ala Ala Lys Tyr Gly Val Gly Thr Pro Ala

420

425

430

Ala Ala Ala Ala Lys Ala Ala Ala Lys Ala Ala Gln Phe Gly Leu Val

435

440

445

Pro Gly Val Gly Val Ala Pro Gly Val Gly Val Ala Pro Gly Val Gly

450

455

460

Val Ala Pro Gly Val Gly Leu Ala Pro Gly Val Gly Val Ala Pro Gly

465

470

475

480

T09240" B08E460

Val Gly Val Ala Pro Gly Val Gly Val Ala Pro Gly Ile Gly Pro Gly  
485 490 495

Gly Val Ala Ala Ala Ala Lys Ser Ala Ala Lys Val Ala Ala Lys Ala  
500 505 510

Gln Leu Arg Ala Ala Ala Gly Leu Gly Ala Gly Ile Pro Gly Leu Gly  
515 520 525

Val Gly Val Gly Val Pro Gly Leu Gly Val Gly Ala Gly Val Pro Gly  
530 535 540

Leu Gly Val Gly Ala Gly Cys Ser Gly Phe Arg Cys Trp Arg Gly Arg  
545 550 555 560

Arg Cys Thr Ser Phe Pro Val Ser Arg Thr Ala  
565 570

<210> 8

<211> 9

<212> PRT

<213> Homo sapiens

<400> 8

Lys Ala Pro Gly Val Gly Gly Ala Phe

1

5

094433 042601  
109240 037626

<210> 9

<211> 7

<212> PRT

<213> Homo sapiens

<400> 9

Arg Ala Ala Ala Gly Leu Gly

1 5

<210> 10

<211> 11

<212> PRT

<213> Homo sapiens

<400> 10

Arg Ser Leu Ser Pro Glu Leu Arg Glu Gly Asp

1 5 10

<210> 11

<211> 9

<212> PRT

<213> Homo sapiens

<400> 11

Lys Ala Ala Lys Ala Gly Ala Gly Leu

1 5

09743818 042601  
T09240" BTBCH.60

<210> 12

<211> 9

<212> PRT

<213> Homo sapiens

<400> 12

Lys Ala Gly Ala Gly Leu Gly Gly Val

1

5

<210> 13

<211> 13

<212> PRT

<213> Homo sapiens

<400> 13

Ala Leu Ala Ala Ala Lys Ala Ala Lys Tyr Gly Ala Ala

1

5

10

<210> 14

<211> 11

<212> PRT

<213> Homo sapiens

<400> 14

09431-04260  
109240 " BTB4.60

Lys Ala Ala Gln Phe Gly Leu Val Pro Gly Val

1

5

10

<210> 15

<211> 11

<212> PRT

<213> Homo sapiens

<400> 15

Lys Ser Ala Ala Lys Val Ala Ala Lys Ala Gln

1

5

10

<210> 16

<211> 9

<212> PRT

<213> Homo sapiens

<400> 16

Arg Ser Leu Ser Pro Glu Leu Arg Glu

1

5

<210> 17

<211> 8

<212> PRT

<213> Homo sapiens

0094331304604



<213> Homo sapiens

<400> 20

Leu Gln Leu Arg Ala Ala Ala Gly

1 5

<210> 21

<211> 8

<212> PRT

<213> Homo sapiens

<400> 21

Ala Asn Leu Arg Ala Ala Ala Gly

1 5

<210> 22

<211> 8

<212> PRT

<213> Homo sapiens

<400> 22

Ala Gly Leu Arg Ala Ala Ala Gly

1 5

<210> 23

<211> 8

091438.042601  
T09240" B7B74.60



<212> PRT

<213> Homo sapiens

<400> 23

Ala Val Leu Arg Ala Ala Ala Gly

1 5

<210> 24

<211> 8

<212> PRT

<213> Homo sapiens

<400> 24

Ala Ser Leu Arg Ala Ala Ala Gly

1 5

<210> 25

<211> 8

<212> PRT

<213> Homo sapiens

<400> 25

Ala Gln Gly Arg Ala Ala Ala Gly

1 5

"09240" 8T884.60

<210> 26

<211> 8

<212> PRT

<213> Homo sapiens

<400> 26

Ala Gln Val Arg Ala Ala Ala Gly

1

5

<210> 27

<211> 8

<212> PRT

<213> Homo sapiens

<400> 27

Ala Gln Ile Arg Ala Ala Ala Gly

1

5

<210> 28

<211> 8

<212> PRT

<213> Homo sapiens

<400> 28

Ala Gln Ala Arg Ala Ala Ala Gly

1

5

004331304601  
"03240"

<210> 29

<211> 8

<212> PRT

<213> Homo sapiens

<400> 29

Ala Gln Leu Arg Gly Ala Ala Gly

1

5

<210> 30

<211> 8

<212> PRT

<213> Homo sapiens

<400> 30

Ala Gln Leu Arg Val Ala Ala Gly

1

5

<210> 31

<211> 8

<212> PRT

<213> Homo sapiens

<400> 31

Ala Gln Leu Arg Ile Ala Ala Gly

00431-04260  
"BTB40" 2240

1

5

<210> 32

<211> 8

<212> PRT

<213> Homo sapiens

<400> 32

Ala Gln Leu Arg Leu Ala Ala Gly

1

5

<210> 33

<211> 8

<212> PRT

<213> Homo sapiens

<400> 33

Ala Gln Leu Arg Ala Gly Ala Gly

1

5

<210> 34

<211> 8

<212> PRT

<213> Homo sapiens

<400> 34

00943318.042604

Ala Gln Leu Arg Ala Val Ala Gly

1

5

<210> 35

<211> 8

<212> PRT

<213> Homo sapiens

<400> 35

Ala Gln Leu Arg Ala Ile Ala Gly

1

5

<210> 36

<211> 8

<212> PRT

<213> Homo sapiens

<400> 36

Ala Gln Leu Arg Ala Leu Ala Gly

1

5

<210> 37

<211> 8

<212> PRT

<213> Homo sapiens

0943818 042604

<400> 37

Ala Gln Leu Arg Ala Ala Gly Gly

1 5

<210> 38

<211> 8

<212> PRT

<213> Homo sapiens

<400> 38

Ala Gln Leu Arg Ala Ala Val Gly

1 5

<210> 39

<211> 8

<212> PRT

<213> Homo sapiens

<400> 39

Ala Gln Leu Arg Ala Ala Ile Gly

1 5

<210> 40

<211> 8

<212> PRT

0943343.042504  
"09240" 8T8E460

<213> Homo sapiens

<400> 40

Ala Gln Leu Arg Ala Ala Leu Gly

1 5

<210> 41

<211> 8

<212> PRT

<213> Homo sapiens

<400> 41

Ala Gln Leu Arg Ala Ala Ala Ala

1 5

<210> 42

<211> 8

<212> PRT

<213> Homo sapiens

<400> 42

Ala Gln Leu Arg Ala Ala Ala Ile

1 5

<210> 43

09743343 042601  
"09240" 8733460

[illegible]

**B**

[illegible][illegible][illegible][illegible][illegible][illegible]

**SECRET**

[illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible]

**B**

[illegible][illegible]



<210> 46

<211> 8

<212> PRT

<213> Homo sapiens

<400> 46

Gly Pro Gly Ala Leu Ala Ala Ala

1 5

<210> 47

<211> 8

<212> PRT

<213> Homo sapiens

<400> 47

Ile Pro Gly Ala Leu Ala Ala Ala

1 5

<210> 48

<211> 8

<212> PRT

<213> Homo sapiens

<400> 48

Leu Pro Gly Ala Leu Ala Ala Ala

1 5

09743813 042604  
T09240 " BTB460



1

5

<210> 52

<211> 8

<212> PRT

<213> Homo sapiens

<400> 52

Val Pro Leu Ala Leu Ala Ala Ala

1

5

<210> 53

<211> 8

<212> PRT

<213> Homo sapiens

<400> 53

Val Pro Val Ala Leu Ala Ala Ala

1

5

<210> 54

<211> 8

<212> PRT

<213> Homo sapiens

T09240"BT3E4.60

<400> 54

Val Pro Gly Ala Gly Ala Ala Ala

1 5

<210> 55

<211> 8

<212> PRT

<213> Homo sapiens

<400> 55

Val Pro Gly Ala Ile Ala Ala Ala

1 5

<210> 56

<211> 8

<212> PRT

<213> Homo sapiens

<400> 56

Val Pro Gly Ala Ala Ala Ala Ala

1 5

<210> 57

<211> 8

<212> PRT

<213> Homo sapiens

09743818.042601

<400> 57

Val Pro Gly Ala Val Ala Ala Ala

1 5

<210> 58

<211> 8

<212> PRT

<213> Homo sapiens

<400> 58

Val Pro Gly Ala Leu Gly Ala Ala

1 5

<210> 59

<211> 8

<212> PRT

<213> Homo sapiens

<400> 59

Val Pro Gly Ala Leu Ile Ala Ala

1 5

<210> 60

<211> 8

0943310420  
"BTB40"

<212> PRT

<213> Homo sapiens

<400> 60

Val Pro Gly Ala Leu Leu Ala Ala

1

5

<210> 61

<211> 8

<212> PRT

<213> Homo sapiens

<400> 61

Val Pro Gly Ala Leu Val Ala Ala

1

5

<210> 62

<211> 8

<212> PRT

<213> Homo sapiens

<400> 62

Val Pro Gly Ala Leu Ala Gly Ala

1

5

<210> 63

094334.04604

<211> 8

<212> PRT

<213> Homo sapiens

<400> 63

Val Pro Gly Ala Leu Ala Ile Ala

1 5

<210> 64

<211> 8

<212> PRT

<213> Homo sapiens

<400> 64

Val Pro Gly Ala Leu Ala Leu Ala

1 5

<210> 65

<211> 8

<212> PRT

<213> Homo sapiens

<400> 65

Val Pro Gly Ala Leu Ala Val Ala

1 5

"09240" BTBETH.60

<210> 66

<211> 8

<212> PRT

<213> Homo sapiens

<400> 66

Val Pro Gly Ala Leu Ala Ala Ala

1

5

<210> 67

<211> 8

<212> PRT

<213> Homo sapiens

<400> 67

Val Pro Gly Ala Leu Ala Ala Gly

1

5

<210> 68

<211> 8

<212> PRT

<213> Homo sapiens

<400> 68

Val Pro Gly Ala Leu Ala Ala Ile

1

5

T09240"STECH/60



<210> 69

<211> 8

<212> PRT

<213> Homo sapiens

<400> 69

Val Pro Gly Ala Leu Ala Ala Leu

1

5

<210> 70

<211> 8

<212> PRT

<213> Homo sapiens

<400> 70

Val Pro Gly Ala Leu Ala Ala Val

1

5

<210> 71

<211> 515

<212> PRT

<213> Homo sapiens

<400> 71

T03240" STB4.60

Gly Gly Val Pro Gly Ala Ile Pro Gly Gly Val Pro Gly Gly Val Phe  
 1 5 10 15  
 Tyr Pro Gly Ala Gly Leu Gly Ala Leu Gly Gly Gly Ala Leu Gly Pro  
 20 25 30  
 Gly Gly Lys Pro Leu Lys Pro Val Pro Gly Gly Leu Ala Gly Ala Gly  
 35 40 45  
 Leu Gly Ala Gly Leu Gly Ala Phe Pro Ala Val Thr Phe Pro Gly Ala  
 50 55 60  
 Leu Val Pro Gly Gly Val Ala Asp Ala Ala Ala Tyr Lys Ala Ala  
 65 70 75 80  
 Lys Ala Gly Ala Gly Leu Gly Gly Val Pro Gly Val Gly Gly Leu Gly  
 85 90 95  
 Val Ser Ala Gly Ala Val Val Pro Gln Pro Gly Ala Gly Val Lys Pro  
 100 105 110  
 Gly Lys Val Pro Gly Val Gly Leu Pro Gly Val Tyr Pro Gly Gly Val  
 115 120 125  
 Leu Pro Gly Ala Arg Phe Pro Gly Val Gly Val Leu Pro Gly Val Pro  
 130 135 140  
 Thr Gly Ala Gly Val Lys Pro Lys Ala Pro Gly Val Gly Gly Ala Phe  
 145 150 155 160

0943818 042601

094381 04601  
T09240" BTCH 460

Ala Gly Ile Pro Gly Val Gly Pro Phe Gly Gly Pro Gln Pro Gly Val  
165 170 175

Pro Leu Gly Tyr Pro Ile Lys Ala Pro Lys Leu Pro Gly Gly Tyr Gly  
180 185 190

Leu Pro Tyr Thr Thr Gly Lys Leu Pro Tyr Gly Tyr Gly Pro Gly Gly  
195 200 205

Val Ala Gly Ala Ala Gly Lys Ala Gly Tyr Pro Thr Gly Thr Gly Val  
210 215 220

Gly Pro Gln Ala Ala Ala Ala Ala Ala Lys Ala Ala Ala Lys Phe  
225 230 235 240

Gly Ala Gly Ala Ala Gly Val Leu Pro Gly Val Gly Gly Ala Gly Val  
245 250 255

Pro Gly Val Pro Gly Ala Ile Pro Gly Ile Gly Gly Ile Ala Gly Val  
260 265 270

Gly Thr Pro Ala Ala Ala Ala Ala Ala Ala Ala Lys Ala Ala  
275 280 285

Lys Tyr Gly Ala Ala Ala Gly Leu Val Pro Gly Gly Pro Gly Phe Gly  
290 295 300

Pro Gly Val Val Gly Val Pro Gly Ala Gly Val Pro Gly Val Gly Val  
305 310 315 320

Pro Gly Ala Gly Ile Pro Val Val Pro Gly Ala Gly Ile Pro Gly Ala  
325 330 335

Ala Val Pro Gly Val Val Ser Pro Glu Ala Ala Ala Lys Ala Ala Ala  
340 345 350

Lys Ala Ala Lys Tyr Gly Ala Arg Pro Gly Val Gly Val Gly Gly Ile  
355 360 365

Pro Thr Tyr Gly Val Gly Ala Gly Gly Phe Pro Gly Phe Gly Val Gly  
370 375 380

Val Gly Gly Ile Pro Gly Val Ala Gly Val Pro Ser Val Gly Gly Val  
385 390 395 400

Pro Gly Val Gly Gly Val Pro Gly Val Gly Ile Ser Pro Glu Ala Gln  
405 410 415

Ala Ala Ala Ala Ala Lys Ala Ala Lys Tyr Gly Val Gly Thr Pro Ala  
420 425 430

Ala Ala Ala Ala Lys Ala Ala Ala Lys Ala Ala Gln Phe Gly Leu Val  
435 440 445

Pro Gly Val Gly Val Ala Pro Gly Val Gly Val Ala Pro Gly Val Gly  
450 455 460

094331042601  
"09240" BTBCH260

Val Ala Pro Gly Val Gly Leu Ala Pro Gly Val Gly Val Ala Pro Gly  
465 470 475 480

Val Gly Val Ala Pro Gly Val Gly Val Ala Pro Gly Ile Gly Pro Gly  
485 490 495

Gly Val Ala Ala Ala Ala Lys Ser Ala Ala Lys Val Ala Ala Lys Ala  
500 505 510

Gln Leu Arg  
515

<210> 72

<211> 49

<212> PRT

<213> Homo sapiens

<400> 72

Ala Ala Ala Gly Leu Gly Ala Gly Ile Pro Gly Leu Gly Val Gly Val  
1 5 10 15

Gly Val Pro Gly Leu Gly Val Gly Ala Gly Val Pro Gly Leu Gly Val  
20 25 30

Gly Ala Gly Val Pro Gly Phe Gly Ala Gly Ala Asp Glu Gly Val Arg  
35 40 45

09/43818.042601  
T09240"8T8E4260

Arg

<210> 73

<211> 171

<212> PRT

<213> Homo sapiens

<400> 73

Gly Val Arg Arg Ser Leu Ser Pro Glu Leu Arg Glu Gly Asp Pro Ser

1

5

10

15

Ser Ser Gln His Leu Pro Ser Thr Pro Ser Ser Pro Arg Val Pro Gly

20

25

30

Ala Leu Ala Ala Ala Lys Ala Ala Lys Tyr Gly Ala Ala Val Pro Gly

35

40

45

Val Leu Gly Gly Leu Gly Ala Leu Gly Gly Val Gly Ile Pro Gly Gly

50

55

60

Val Val Gly Ala Gly Pro Ala Ala Ala Ala Ala Ala Lys Ala Ala

65

70

75

80

Ala Lys Ala Ala Gln Phe Gly Leu Val Gly Ala Ala Gly Leu Gly Gly

85

90

95

Leu Gly Val Gly Gly Leu Gly Val Pro Gly Val Gly Gly Leu Gly Gly  
100 105 110

Ile Pro Pro Ala Ala Ala Ala Lys Ala Ala Lys Tyr Gly Ala Ala Gly  
115 120 125

Leu Gly Gly Val Leu Gly Gly Ala Gly Gln Phe Pro Leu Gly Gly Val  
130 135 140

Ala Ala Arg Pro Gly Phe Gly Leu Ser Pro Ile Phe Pro Gly Gly Ala  
145 150 155 160

Cys Leu Gly Lys Ala Cys Gly Arg Lys Arg Lys  
165 170

<210> 74

<211> 183

<212> PRT

<213> Homo sapiens

<400> 74

Ala Ala Ala Gly Leu Gly Ala Gly Ile Pro Gly Leu Gly Val Gly Val  
1 5 10 15

Gly Val Pro Gly Leu Gly Val Gly Ala Gly Val Pro Gly Leu Gly Val  
20 25 30

094331.04601  
T09240" BTB4.60

Gly Ala Gly Val Pro Gly Phe Gly Ala Val Pro Gly Ala Leu Ala Ala

35

40

45

Ala Lys Ala Ala Lys Tyr Gly Ala Ala Val Pro Gly Val Leu Gly Gly

50

55

60

Leu Gly Ala Leu Gly Gly Val Gly Ile Pro Gly Gly Val Val Gly Ala

65

70

75

80

Gly Pro Ala Ala Ala Ala Ala Ala Ala Lys Ala Ala Ala Lys Ala Ala

85

90

95

Gln Phe Gly Leu Val Gly Ala Ala Gly Leu Gly Gly Leu Gly Val Gly

100

105

110

Gly Leu Gly Val Pro Gly Val Gly Gly Leu Gly Gly Ile Pro Pro Ala

115

120

125

Ala Ala Ala Lys Ala Ala Lys Tyr Gly Ala Ala Gly Leu Gly Gly Val

130

135

140

Leu Gly Gly Ala Gly Gln Phe Pro Leu Gly Gly Val Ala Ala Arg Pro

145

150

155

160

Gly Phe Gly Leu Ser Pro Ile Phe Pro Gly Gly Ala Cys Leu Gly Lys

165

170

175

Ala Cys Gly Arg Lys Arg Lys

180

09743318-042601